

Gastric bypass alters sweet taste function

November 2 2010

Gastric bypass surgery decreases the preference for sweet-tasting substances in obese rats, a study finding that could help in developing safer treatments for the morbidly obese, according to Penn State College of Medicine researchers.

"Roux-en-Y [gastric bypass surgery](#) is the most common effective treatment for [morbid obesity](#)," said Andras Hajnal, M.D., Ph.D., associate professor, Department of Neural and Behavioral Science and Surgery. "Many patients report altered taste preferences after having the procedure."

This surgery involves the creation of a small gastric pouch and bypassing a portion of the upper [small intestine](#). Unlike other weight-reduction methods, it produces substantial and durable weight loss and significant improvements in obesity-related medical conditions including diabetes.

Study results in obese rats suggest that post-surgery changes in the gastrointestinal anatomy affect change in the brain that relate to taste.

Obese rats given gastric bypass surgery showed a reduced preference for high concentration sucrose water when compared to obese rats that did not have surgery. Researchers observed a similar decrease in preference with other sweet-tasting substances, but not for salty, sour or bitter substances. Researchers observed no change in preference in lean rats that had gastric bypass surgery.

The obese rats used do not have the ability to produce the receptor for

feeling satiated shortly after a meal because they lack the gut hormone CCK-1. As a result, these rats consumed larger meals and, over time, became obese and developed type-2 diabetes. Interestingly, previous studies lead by the Penn State investigators found an increased sweet preference in these rats, which is also often seen in people struggling with weight management.

"It appears that an uncontrolled appetite may get further boost from altered taste functions during development of obesity and diabetes," Hajnal said. "How much of this vicious circle is due to changes in the neurons inside the brain, which receive taste sensations from the tongue and report to the higher order motivational brain centers, we don't know."

The researchers recorded the activity of 170 taste-responsive neurons in the brain. These showed a shift in the neurons' firing activity similar to the behavioral response, which was measured in lick rates of the rats within a ten-second time period. Neurons in the obese rats' brain responded more vigorously to higher-concentration sucrose water placed on the tongue when compared to lean rats. These effects were reversed by gastric bypass surgery and matched the response of lean rats -- a preference for lower concentration sucrose.

The rats that had gastric bypass surgery lost weight comparable to humans who received the surgery -- 26 to 30 percent of their weight -- and maintained the loss for a long period of time after surgery. Following surgery, the obese rats also showed a higher tolerance for glucose, indicating improvement in diabetes.

"This supports the applicability of this rat model of Roux-en-Y gastric bypass to humans and also suggests that the observed taste changes following the surgery were not related to 'human factors' such as awareness and compliance to dietary and behavioral interventions,"

Hajnal said.

The researchers published their findings in the October issue of the *"American Journal of Physiology, Gastrointestinal and Liver Physiology."* The National Institute of Diabetes and Digestive and Kidney Diseases and the National Institute of General Medical Sciences funded this work.

"These findings confirm obesity-related alterations in taste functions and demonstrate the ability of gastric bypass surgery to alleviate these modifications," Hajnal said. "We do not suggest, however, that the findings reported in this paper are the only neural consequences of gastric bypass surgery related to altered postsurgical food preferences. Nevertheless, understanding the underlying mechanisms by which gastric bypass surgery affects taste may help in identifying therapeutic targets that mimic the beneficial effects of the surgery on appetite control and food choices, without the risks and complications of an invasive surgical procedure."

Provided by Pennsylvania State University

Citation: Gastric bypass alters sweet taste function (2010, November 2) retrieved 1 May 2024 from <https://medicalxpress.com/news/2010-11-gastric-bypass-sweet-function.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--