

# Study examines risk of heterosexual HIV transmission in China

November 8 2010

---

A new study by researchers at the Johns Hopkins Bloomberg School of Public Health examines the burden of HIV and sexually transmitted disease among male clients of the commercial sex industry in China's Sichuan province. Since 2007, heterosexual transmission has replaced injecting drug use as the primary transmission mode of all HIV infections in China. The study is available online in advance of publication in the journal *Sexually Transmitted Infections*.

"We found a large burden of syphilis infection coupled with high-risk sexual and substance use behaviors among male clients of [sex workers](#)," said Cui Yang, lead author of the study and a postdoctoral fellow with Bloomberg School's Department of Epidemiology. "Without effective interventions to prevent [HIV transmission](#) to the male client population, the current epidemic may expand into the general population."

For the study, Yang and her colleagues examined the prevalence of HIV, syphilis and [hepatitis C virus](#) (HCV) in a sample of 600 male clients of female sex workers in three urban areas in Sichuan province, [China](#). Blood test results indicated that HIV prevalence was 1.5 percent; 5.3 percent were positive for syphilis and 8.7 percent positive for HCV. The overall prevalence of consistent condom use with female sex workers was 30 percent. The study also found a significant correlation between syphilis infection among male clients and not having local household registration, snorting heroin and washing genitals after having sex with a female sex worker.

"Our data raises concerns about the future heterosexual transmission of HIV to the general population in China since unprotected sex with female sex workers has become the norm," said Kenrad Nelson, MD, senior author of the study and professor in the Department of Epidemiology at the Bloomberg School of Public Health.

Authors of "HIV, syphilis, [hepatitis C](#) and risk behaviours among commercial sex male clients in Sichuan province, China" include Carl Latkin and Cunling Wang of the Bloomberg School of Public Health, and Rongsheng Luan of the West China School of Public Health at Sichuan University.

Provided by Johns Hopkins University Bloomberg School of Public Health

Citation: Study examines risk of heterosexual HIV transmission in China (2010, November 8) retrieved 9 April 2024 from <https://medicalxpress.com/news/2010-11-heterosexual-hiv-transmission-china.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--