

Side effects of hormonal breast cancer therapy increased; may affect treatment adherence

November 9 2010

Women being treated for breast cancer with aromatase inhibitors may experience extremely low estrogen levels resulting in a wide variety of side effects that a typical postmenopausal woman without cancer may not experience.

Data presented at the Ninth Annual AACR Frontiers in [Cancer Prevention](#) Conference, held here Nov. 7-10, 2010, showed that women assigned to take aromatase inhibitors had increases in side effects such as hot flashes, decreased appetite, fatigue, fever, breast sensitivity, etc.

"Aromastase inhibitors represent one of the most major advances in breast cancer treatment," said Lisa Gallicchio, Ph.D., an epidemiologist at The Prevention and Research Center at Mercy Medical Center, Baltimore. "Their incorporation into the breast cancer treatment armamentarium has led to impressive reductions in breast cancer recurrence and mortality rates.

"Despite this, many breast cancer patients stop taking their aromatase inhibitor treatment—which is usually prescribed for five years—or do not adhere to their treatment prescription," she said, adding that this may be due to, at least in part, the side effects associated with the drugs.

To better define the full spectrum of side effects associated with aromatase inhibitor treatment, Gallicchio and colleagues surveyed 100

women with breast cancer who were about to start treatment with aromatase inhibitors. Their side effects were compared with those of 200 similarly-aged women without a history of breast cancer.

Questionnaires about symptoms were completed prior to treatment initiation by women with breast cancer and at the start of the study for the healthy women. Women were followed for six months and completed additional questionnaires at three months and at the completion of the study.

Women taking [aromatase inhibitors](#) were five times more likely to report having hot flashes, breast sensitivity and chest pain than healthy women. In addition, they were four times more likely to report night sweats, cold sweats and hair loss and about three times more likely to report leg cramps, weight gain, sleep disturbance, tendency to take naps and forgetfulness. Other increased symptoms included intestinal gas, cough, depression, interrupted sleep and irritability.

"We know that aromatase [inhibitors](#) are effective in treating [breast cancer](#)," Gallicchio said. "Knowing the side effects of aromatase inhibitor treatment and how to treat them is critical for keeping women on their aromatase inhibitor treatment and improving their chances of surviving and living cancer free."

Provided by American Association for Cancer Research

Citation: Side effects of hormonal breast cancer therapy increased; may affect treatment adherence (2010, November 9) retrieved 1 May 2024 from <https://medicalxpress.com/news/2010-11-side-effects-hormonal-breast-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
