

# What happens after traumatic brain injury occurs?

November 1 2010

---

Results from a study published in the November issue of *The Journal of Nuclear Medicine* (JNM) show that powerful imaging techniques—positron emission tomography (PET) fused with magnetic resonance imaging (MRI)—are helping researchers better understand the long-term functional and structural changes that take place after traumatic brain injury (TBI).

"The results provide new insights into the progressive nature of the brain changes occurring following TBI, demonstrating that changes in brain structure and function after TBI are dynamic and continue to progress and evolve for months," said Viviane Bouilleret, MD, PHD, Univ Paris-Sud, one of the authors of the study, "Progressive Metabolic and Structural Cerebral Perturbations After [Traumatic Brain Injury](#): An In Vivo Imaging Study in the Rat."

The paper reports the first study to use repeated PET and MRI of live animals to characterize the long-term progressive functional and structural brain changes that occur following experimental TBI. The model used is the lateral fluid percussion injury model in rats, the most widely utilized and validated animal model in basic TBI research. The images were assessed using a variety of sophisticated, state-of-the-art image analysis approaches.

Widespread decreases in brain functioning were seen in specific brain regions, many of which are remote from the site of direct trauma and unaccompanied by signs of injury on the MRI. The hippocampus, a

brain structure critical to memory and emotion, is the key area of these changes. This may have implications for the pathophysiology of some of the long-term neurological and psychiatric morbidity, seen following TBI, even when abnormalities are not obvious on structural MRI. The severity of the neurological deficits and changes on MRI soon after the injury is predictive of severity of long-term neurodegenerative changes.

Traumatic [brain injury](#) is a leading cause of death and disability in children and young adults around the world. Long-term disabilities following TBI include: cognitive disabilities such as disorders in attention, memory, learning and executive functioning; psychiatric and behavioral problems including depression, anxiety, aggression, agitation and social inappropriateness; and epilepsy. Currently there are no effective interventions to reduce the incidence or severity of these problems.

The findings of this study indicate that secondary structural and functional changes in the brain continue to occur for many months post-TBI, which has implications for understanding the mechanisms underlying the long-term neurological and psychiatric consequences. Importantly, these findings indicate that there is a time window during which intervention could modulate these processes and mitigate against these disabling consequences of TBI.

This study has also provided a practical paradigm for testing the biological effectiveness of potential therapies prior to embarking on expensive and lengthy clinical trials. "The approach would also be applicable to the study of other neurological diseases, such as stroke, dementia, multiple sclerosis brain infections and epilepsy, which are associated with long-term progressive degenerative changes in the brain," Bouilleret said.

**More information:** "Progressive Metabolic and Structural Cerebral

Perturbations After Traumatic Brain Injury: An In Vivo Imaging Study in the Rat," [jnm.snmjournals.org/](http://jnm.snmjournals.org/)

Provided by Society of Nuclear Medicine

Citation: What happens after traumatic brain injury occurs? (2010, November 1) retrieved 12 September 2024 from <https://medicalxpress.com/news/2010-11-traumatic-brain-injury.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.