

# Women with personal history of breast cancer should be screened with MRI

November 29 2010

---

Women with a personal history of breast cancer should consider annual screening with MRI in addition to mammography, according to a study presented today at the annual meeting of the Radiological Society of North America (RSNA).

The American Cancer Society (ACS) guidelines currently recommend annual screening with breast MRI in [women](#) with a known gene mutation or with a strong [family history](#) indicating a lifetime risk of [breast cancer](#) greater than 20 percent. However, the guidelines say there is insufficient evidence to recommend for or against MRI screening in women who have already had breast cancer themselves.

"In our study using breast MRI screening, we actually detected proportionally more cancers in women with a personal history of breast cancer, compared with those women with a genetic mutation or strong family history who are currently recommended to have breast MRI," said Wendy B. DeMartini, M.D., assistant professor in the Department of Radiology at the University of Washington Medical Center and Seattle Cancer Care Alliance in Seattle. "Further, women with a personal history were less likely to be recalled for additional testing and less likely to have a biopsy for a false positive MRI finding."

Dr. DeMartini and colleagues performed a retrospective review of initial screening breast MRI examinations of 1,026 women from January 2004 to June 2009. Of the 1,026 women, 327 had a genetic or family history of breast cancer and 646 had a personal history of treated breast cancer.

Overall, MRI testing identified 25 of 27 cancers in the group for a sensitivity rate of 92.6 percent.

The cancer yield in the women with a personal history of breast cancer (3.1 percent) was double that of the women with a genetic or family history (1.5 percent). Specificity in women with a personal history was 93.6 percent, compared with 86.3 percent for the other group.

Specificity refers to the accuracy of the exam in correctly ruling out cancer where it is not present, resulting in lower recall and biopsy rates due to false-positive findings.

Biopsy was recommended in 9.3 percent of the women with a personal history of breast cancer, compared with 15 percent of the genetic and family history group. The positive predictive value of biopsy was also higher in the personal history group, with 35.7 percent of biopsies yielding cancer, compared with only 12.2 percent in the other group.

"Our findings show that the diagnostic performance of MRI in patients with a personal history of treated breast cancer supports consideration of screening MRI as an adjunct to mammography," Dr. DeMartini said.

"Additional studies such as ours are necessary to establish guidelines for screening this important group of women."

Provided by Radiological Society of North America

Citation: Women with personal history of breast cancer should be screened with MRI (2010, November 29) retrieved 10 April 2024 from <https://medicalxpress.com/news/2010-11-women-personal-history-breast-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--