

Buprenorphine treatment produces improved outcome for babies born addicted

December 9 2010

Babies born into the world addicted to drugs because of their mother's dependence on pain medication, or opioids, may be weaned off the substance more comfortably, with a shorter hospital stay and at a reduced cost, if the mother receives a new treatment option during pregnancy.

A [New England Journal of Medicine](#) study published Dec. 9, co-authored by Vanderbilt's Peter Martin, M.D., director of the Division of Addiction Psychiatry, found that the newer buprenorphine is at least as good for both mother and child as the standard care [methadone](#), when both were combined with comprehensive care treatment of opioid dependence in [pregnant women](#).

"The baby comes into the world addicted to what the mother was taking," Martin said.

"And everyone who comes in touch with these unfortunate [babies](#) realizes that they are very uncomfortable and incessantly restless.

"From this study we can say that both the mothers and the babies did equally well taking buprenorphine or methadone.

"However, we demonstrated a statistically significant improvement above the standard of care in important outcomes in the babies of mothers who received buprenorphine during the pregnancy compared to those who were administered methadone," he said.

The eight site, international, double-blind, double-dummy, flexible-dosing, randomized controlled trial is part of the MOTHER project, which stands for Maternal Opioid Treatment: Human Experimental Research.

Buprenorphine had previously not been well studied in pregnancy, although it is now widely prescribed to treat opioid addiction.

"It was only a matter of time before women who were receiving buprenorphine while they become pregnant were faced with the decision to continue it or change to methadone, the standard of care," Martin said.

"For this reason, it was considered very important to determine whether buprenorphine was at least as safe as methadone, so that doctors would know how to advise their patients."

Study results showed that babies of mothers who received [buprenorphine](#) compared to those who received methadone throughout pregnancy needed significantly less morphine to treat their neonatal abstinence syndrome, had shorter hospital stays (10 days vs. 17.5 days), and shorter duration of treatment for neonatal abstinence syndrome (4.1 days vs. 9.9 days).

The authors state that improved treatment options should reduce the public health and medical costs associated with the treatment of neonates exposed to opioids, estimated at \$70.6 million to \$112.6 million in 2009 in the United States.

Provided by Vanderbilt University Medical Center

Citation: Buprenorphine treatment produces improved outcome for babies born addicted (2010,

December 9) retrieved 11 May 2024 from

<https://medicalxpress.com/news/2010-12-buprenorphine-treatment-outcome-babies-born.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.