

Impaired clearance, not overproduction of toxic proteins, may underlie Alzheimer's disease

December 9 2010

In Alzheimer's disease, a protein fragment called beta-amyloid accumulates at abnormally high levels in the brain. Now researchers funded by the National Institutes of Health have found that in the most common, late-onset form of Alzheimer's disease, beta-amyloid is produced in the brain at a normal rate but is not cleared, or removed from the brain, efficiently. In addition to improving the understanding of what pathways are most important in development of Alzheimer's pathology, these findings may one day lead to improved biomarker measures for early diagnosis as well as a new approach to treating this devastating disorder.

Many believe that accumulation of abnormal levels of beta-amyloid in the <u>brain</u> initiates a cascade of events leading to the death of <u>brain cells</u> and ultimately to <u>dementia</u>. In the rare, early-onset forms of Alzheimer's that are linked to <u>genetic mutations</u> there is a marked increase in betaamyloid production. In the more common, late-onset form of Alzheimer's, the mechanisms leading to increased beta-amyloid levels are not well understood.

The study, published in Science, was led by senior author Randall Bateman, M.D., an assistant professor of neurology at Washington University in St. Louis. Dr. Bateman and his colleagues previously reported an innovative procedure to measure beta-amyloid levels over time in cerebrospinal fluid (CSF) – the fluid that bathes the brain. In the



new study, the researchers used that procedure to measure beta-amyloid production and clearance rates in study volunteers with Alzheimer's disease and in age-matched volunteers free of the disease.

"These findings may help point us toward better diagnostic tests and effective therapies. The next question is what is causing the decreased clearance rate," said Dr. Bateman.

Prior studies in animals suggest several possible explanations for the slower clearance of beta-amyloid in late-onset Alzheimer's. One possibility is that as beta-amyloid accumulates, it acts as a sink for more of the protein, trapping it within the brain. The researchers believe that sorting out these mechanisms is likely to help speed the development of new drugs for the disease.

"Abnormal protein deposits within the brain are a hallmark not only of Alzheimer's disease, but of many neurological disorders. With knowledge about how these proteins accumulate, we may be able to slow that process and reduce the damage to the brain," said Roderick Corriveau, Ph.D., a program director at NIH's National Institute of Neurological Disorders and Stroke (NINDS).

Dr. Bateman and his colleagues measured beta-amyloid levels in the CSF of 12 patients with late-onset Alzheimer's and 12 cognitively normal individuals. The participants' average age was 74. Their CSF was sampled every hour for 36 hours, via a one-time lumbar puncture. Throughout the procedure, they were encouraged to stay in bed.

During the first nine hours, the participants received an IV drip containing leucine, a protein building block, which had been labeled with a non-radioactive isotope. By measuring the amount of labeled betaamyloid by mass spectroscopy over time, the researchers were able to calculate how fast it was produced in the participants' brains and how



fast it was cleared. The clearance of beta-amyloid was about 30 percent slower in individuals with <u>Alzheimer's disease</u> than in cognitively normal individuals. This is consistent with previous findings that beta-amyloid is decreased in the CSF of people with Alzheimer's and in those with mild cognitive impairment at high risk of developing the disease.

Dr. Bateman's procedure has enabled researchers to investigate the effects of new drugs aimed at tamping down beta-amyloid levels. The procedure also could be a starting point toward a shorter, simpler test – perhaps a blood test – that measures beta-amyloid clearance rates to detect

Alzheimer's prior to the first symptoms.

"This study is significant in that it reports the first measurement of betaamyloid production and clearance in Alzheimer's," said Marcelle Morrison-Bogorad, Ph.D., director of the Division of Neuroscience at the National Institute on Aging (NIA). "For years scientists believed that it was the overproduction of beta-amyloid that led to its accumulation in the brain. These new findings shift the emphasis to clearance of betaamyloid. This may lead to development of a diagnostic test as well as identification of new therapeutic targets."

More information: Mawuenyega, KG et al. "Decreased clearance of amyloid-beta in the CNS of humans with Alzheimer's disease." *Science*, published online December 9, 2010.

Provided by National Institutes of Health

Citation: Impaired clearance, not overproduction of toxic proteins, may underlie Alzheimer's disease (2010, December 9) retrieved 4 May 2024 from https://medicalxpress.com/news/2010-12-impaired-clearance-overproduction-toxic-proteins.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.