

New test announced for major killer of lung transplant patients

December 30 2010

A lung transplant can mean a new chance at life. But many who receive one develop a debilitating, fatal condition that causes scar tissue to build up in the lungs and chokes off the ability to breathe.

University of Michigan researchers hope a new diagnostic tool they developed to predict bronchiolitis obliterans syndrome (BOS) will allow doctors to intervene earlier and, ultimately, to provide life-saving treatments.

BOS is the leading cause of death for those who survive one year after lung transplantation and more than half of recipients will develop BOS within five years. There is currently no cure.

Vibha Lama, M.D., M.S., an assistant professor of pulmonary and critical care medicine at the University of Michigan Medical School, led a team of U-M researchers who recently discovered that patients who had a high number of [stem cells](#) in their lungs six months after transplantation were much more likely to develop BOS than those with lower counts.

"Our study provides the first indication of the important role these [cells](#) play in both human repair and disease," Lama says. "It's very important from the clinical perspective because we didn't previously have any strong [biomarkers](#) for BOS."

The findings were recently published online ahead of print publication in

the *American Journal of Respiratory and [Critical Care Medicine](#)*.

The translational study also highlights the importance of the lab-to-bedside cooperation of basic and clinical research, Lama says.

While the exact relationship between the mesenchymal stromal cells and BOS remains unclear, doctors know that most of the cells originate with the donor and not the recipient. Spikes in cell counts are seen shortly after transplantation as the body responds to the injury; those levels usually taper off, but a second rise of cell counts after about six months is linked to a patient's likelihood of developing BOS.

In 2007, Lama and her colleagues published another discovery about the stem cells, revealing that the cells reside in the transplanted organs, independent of their more commonly known association with bone marrow. That study led to the further exploration of the cells' involvement with chronic transplant rejection.

The new findings also have the potential to spur research that will help people suffering from other types of lung disease, such as idiopathic pulmonary fibrosis, known as IPF.

Having the biomarker will also allow researchers to readily identify a population of patients ideal for testing new drug interventions and therapies.

"By the time we usually diagnose BOS, there's already been a huge decline in lung function," Lama says. "If we can find the disease early, we can potentially do something about it."

Methodology: [Mesenchymal](#) stromal cells were measured in 405 bronchoalveolar lavage fluid samples obtained from 162 [lung transplant](#) recipients and patients were observed for BOS development.

Provided by University of Michigan Health System

Citation: New test announced for major killer of lung transplant patients (2010, December 30)
retrieved 9 April 2024 from

<https://medicalxpress.com/news/2010-12-major-killer-lung-transplant-patients.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--