

Teens with pelvic inflammatory disease rarely seen in outpatient setting due to costs

January 7 2011, By Ekaterina Pesheva

Hospitalizing teen girls with pelvic inflammatory disease (PID) costs six times as much as treating them in the emergency room, and up to 12 times more than treating them in an outpatient clinic, according to a small study conducted at the Johns Hopkins Children's Center.

The findings, published online in the December issue of [Sexually Transmitted Diseases](#), are based on an analysis of 172 patient visits among 152 girls, 12 to 21 years of age, with PID.

The researchers say the study underscores the need for earlier diagnosis and treatment of these patients not only to help contain costs, but also, more importantly, to prevent PID complications like [chronic pelvic pain](#) and [infertility](#).

“The dollar cost of PID hospitalizations and ER care is important, but at the same time we should take steps to individualize PID care and tailor it to each girl's specific age and circumstances to help her understand how to prevent this from happening again,” says lead investigator Maria Trent, M.D., M.P.H., a Hopkins Children's pediatrician who studies teen sexual and reproductive health.

Of the 135 outpatient visits in the study, only 14 involved visits to a regular clinic, while 121 were visits to the [emergency room](#), probably signaling lack of reliable primary care for many of the teens in the study, the researchers say. Outpatient visits cost on average \$701, compared with \$1,382 for treatment in the emergency room.

By comparison, inpatient treatment in the hospital costs on average \$8,480 per patient per episode. Thirty-seven of the 172 visits resulted in hospitalizations because of severe or advanced PID. Hospital charges were even higher — \$13,360 — for a small subset of girls with PID who required treatment on a psychiatric unit, a finding that suggests how vulnerable sexually active teens with mental health disorders might be.

PID, an inflammation of the reproductive organs, is a complication of untreated sexually transmitted infections like chlamydia and gonorrhea, among other bacterial infections, and affects more than 1 million women in the United States each year, according to the U.S. Centers for Disease Control and Prevention. More than 100,000 of these women develop fertility problems as a result of their infections.

Provided by Johns Hopkins University

Citation: Teens with pelvic inflammatory disease rarely seen in outpatient setting due to costs (2011, January 7) retrieved 18 April 2024 from <https://medicalxpress.com/news/2011-01-teens-pelvic-inflammatory-disease-rarely.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--