

# Risk of cancer increases with exposure to low-dose radiation

February 7 2011

---

Exposure to low-dose radiation from cardiac imaging and other procedures after a heart attack is associated with an increased risk of cancer, found a new study published in *CMAJ (Canadian Medical Association Journal)*.

The use of procedures with low-dose [ionizing radiation](#), such as computed tomography (CT) angiography and nuclear scans, is increasing which has led to mounting concern in the medical community that patients may be at increased risk of cancer. For patients with known or suspected coronary artery disease, the trend towards increased use of these procedures is particularly strong. In many centres, these procedures are replacing those that do not use radiation, such as stress tests on exercise treadmills and echocardiography. However, little is known about the effects of exposure to radiation and the risk of cancer.

The study, conducted by researchers from the McGill University Health Centre (MUHC) and the Jewish General Hospital in Montréal, Quebec, looked at data on 82 861 patients who had a [heart attack](#) between April 1996 and March 2006 but no history of cancer. Of this number, 77% underwent at least one cardiac procedure with low-dose ionizing radiation within one year of the attack.

"We found a relation between the cumulative exposure to low-dose ionizing radiation from cardiac imaging and therapeutic procedures after acute myocardial infarction, and the risk of incident [cancer](#)," writes Dr. Louise Pilote, researcher in epidemiology at the Research Institute of the

MUHC and director of the Division of Internal Medicine at the MUHC with coauthors. "Although most patients were exposed to low or moderate levels of radiation, a substantial group were exposed to high levels and in general tended to be younger male patients with fewer comorbidities."

The median age of patients was 63.2 years and 31.7% were women. Patients whose treating physician was a cardiologist had higher levels of exposure to radiation compared with those whose treating physician was a general practitioner. There were 12 020 incident cancers detected during follow up, with two-thirds of the cancers affecting the abdomen/pelvis and chest areas. "These results call into question whether our current enthusiasm for imaging and therapeutic procedures after acute myocardial infarction should be tempered," conclude the authors. "We should at least consider putting into place a system of prospectively documenting the imaging tests and procedures that each patient undergoes and estimating his or her cumulative exposure to low-dose ionizing radiation."

In a related commentary, Mathew Mercuri, McMaster University, and coauthors write that, although the radiation exposure of many tests is often low, "the frequency with which such tests are performed may pose a population risk." They state the best solution may be prevention, which could mean using procedures with lower or no radiation exposure, especially if there are multiple procedures involved. They call for programs to track radiation doses to help patients and physicians track the risk of cumulative exposure.

Provided by Canadian Medical Association Journal

Citation: Risk of cancer increases with exposure to low-dose radiation (2011, February 7) retrieved 17 April 2024 from

<https://medicalxpress.com/news/2011-02-cancer-exposure-low-dose.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.