

Cell-phone use not related to increased brain cancer risk in new study

February 17 2011

Radio frequency exposure from cell phone use does not appear to increase the risk of developing brain cancers by any significant amount, a study by University of Manchester scientists suggests.

The researchers used publically available data from the UK Office of National Statistics to look at trends in rates of newly diagnosed brain cancers in England between 1998 and 2007.

The study, published in the journal *Bioelectromagnetics*, reported no statistically significant change in the incidence of brain cancers in men or women during the nine-year time period under observation.

"Cell phone use in the United Kingdom and other countries has risen steeply since the early 1990s when the first digital cell phones were introduced," said lead researcher Dr Frank de Vocht, an expert in occupational and environmental health in the University of Manchester's School of Community-Based Medicine.

"There is an on-going controversy about whether <u>radio frequency</u> exposure from cell phones increases the risk of <u>brain cancer</u>. Our findings indicate that a causal link between cell phone use and cancer is unlikely because there is no evidence of any significant increase in the disease since their introduction and rapid proliferation"

The authors say that because there is no plausible <u>biological mechanism</u> for <u>radio waves</u> to damage our <u>genes</u> directly, thereby causing cells to



become cancerous, radio frequency exposure, they argue, if related to cancer is more likely to promote growth in an existing brain tumour.

As such, the researchers say they would expect an increase in the number of diagnosed cases within five to 10 years of the introduction of cell phones and for this increase to continue as cell phone use became more widespread. The 1998 to 2007 study period would therefore relate to the period 1990 to 2002 when cell phone use in the UK increased from zero to 65% of households.

The team, which included researchers from the Institute of Occupational Medicine in Edinburgh and Drexel University, Philadelphia, found a small increase in the incidence of cancers in the temporal lobe of 0.6 cases per 100,000 people or 31 extra cases per year in a population of 52 million. Brain cancers of the parietal lobe, cerebrum and cerebellum in men actually fell slightly between 1998 and 2007.

"Our research suggests that the increased and widespread use of cell phones, which in some studies was associated to increased brain cancer risk, has not led to a noticeable increase in the incidence of brain cancer in England between 1998 and 2007," said Dr de Vocht.

"It is very unlikely that we are at the forefront of a brain cancer epidemic related to cell phones, as some have suggested, although we did observe a small increased rate of brain cancers in the temporal lobe corresponding to the time period when cell phone use rose from zero to 65% of households. However, to put this into perspective, if this specific rise in tumour incidence was caused by <u>cell phone</u> use, it would contribute to less than one additional case per 100,000 population in a decade.

"We cannot exclude the possibility that there are people who are susceptible to radio-frequency exposure or that some rare brain cancers



are associated with it but we interpret our data as not indicating a pressing need to implement public health measures to reduce radio-frequency exposure from cell phones."

Provided by University of Manchester

Citation: Cell-phone use not related to increased brain cancer risk in new study (2011, February 17) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2011-02-cell-phone-brain-cancer.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.