

# Effective treatment of Lyme-disease-related arthritis depends on proper diagnosis

February 2 2011

---

– Early, correct diagnosis is the best way to prevent the development of Lyme arthritis in individuals with the tick-borne illness, according to a paper published in the *Journal of the American Academy of Orthopaedic Surgeons* (JAAOS). In patients who do develop the condition, most cases can be treated successfully with antibiotics, the review found.

"Lyme [arthritis](#) occurs commonly in patients with [Lyme disease](#) and should be considered when evaluating patients with joint complaints and who live in areas where the disease occurs," said study author Aristides Cruz, MD, chief orthopaedic resident, Yale-New Haven Hospital.

"When diagnosed early, most patients do not develop Lyme arthritis. But when correct diagnosis is delayed, arthritis can occur and requires intervention before permanent joint damage develops."

Lyme arthritis can be defined as a painful, swollen joint, that causes a stiffness similar to osteoarthritis, and occurs most commonly in the late stages of Lyme disease, usually several months after the onset of the disease. About 60 percent of patients who are left untreated for Lyme disease in its early stages will develop Lyme arthritis, Dr. Cruz said.

"Lyme arthritis occurs when the spirochete, the bacteria that causes the disease, invades the joints and causes inflammation to the tissue that lines the joints," he said. "If left untreated, this inflammatory response can cause the cartilage within the joints to become damaged."

Dr. Cruz said most cases of Lyme arthritis are brief and involve a single

joint, most typically the knee. The ankle, elbow, hip and wrist may also be affected, and some patients may develop a fever. Following an initial period of joint pain and swelling lasting from a week to several months, symptoms may recur until the disease is diagnosed. In the acute and late stages, Lyme disease may be difficult to distinguish from other disease processes, he added.

"Diagnosing Lyme arthritis can be problematic, especially in patients who do not report a rash and who have fever, since it can be confused with a bacterial joint infection, particularly in children," Dr. Cruz said. "In general, the diagnosis is multifaceted and involves a thorough understanding of the common clinical and epidemiologic features of the disease, along with careful understanding of the tests used in diagnosis."

"In most patients, Lyme arthritis can be successfully treated with oral antibiotics," Dr. Cruz said. "Nonsteroidal anti-inflammatory drugs may also be given to help manage symptoms while the patient is improving." A small number of patients may continue to have symptoms even after treatment with antibiotics, Dr. Cruz said. In these patients, surgical treatment may be needed to remove the diseased synovial tissue.

To help ensure a speedy diagnosis, Dr. Cruz said individuals who live in areas where Lyme disease is known to be present should look for the following symptoms and report them to their physician:

- Rash which develops near an insect bite on the skin
- Aching, redness or swelling in one or more joints especially the knee
- Fever

- Lethargy or malaise
- Headache

Dr. Cruz said that although rash occurs in the majority of patients exposed to the Lyme bacteria, in many patients who develop Lyme arthritis, the rash either never develops or goes unnoticed.

"In the classic form of Lyme disease, a rash occurs, which resembles a bull's eye," he said. "The fact that this rash does not always occur or is not reported in patients with Lyme arthritis makes the condition even more difficult to accurately diagnose."

Patients who suspect they may have been exposed to ticks carrying Lyme disease should speak to their physicians about the tests used to determine if Lyme disease is present, he said.

"Lyme arthritis has an excellent prognosis with appropriate treatment," Dr. Cruz said. "Most patients respond well to a single course of antibiotics. The key is to diagnose Lyme disease early to avoid development of arthritis, and to differentiate between Lyme arthritis and other types of arthritis in symptomatic patients."

More about Lyme disease:

According to the Centers for Disease Control (CDC), about 30,000 cases of Lyme disease were confirmed in 2009. It occurs most commonly in the Northeast, the Great Lakes region of the Midwest and in the Pacific Northwest. To help prevent Lyme disease and associated arthritis, the CDC recommends the following steps:

- Avoid wooded or bushy areas with lots of leaf litter and high

grass

- Take extra precautions during May, June and July when ticks are most active
- Use insect repellent with 20 percent - 30 percent DEET on exposed skin and clothing
- Wear long pants, long sleeves, and long socks to keep ticks off your skin. Tuck pants into boots or socks and wear light-colored clothing so ticks are easier to spot
- Check your skin and clothes for ticks before going indoors

Provided by American Academy of Orthopaedic Surgeons

Citation: Effective treatment of Lyme-disease-related arthritis depends on proper diagnosis (2011, February 2) retrieved 20 April 2024 from

<https://medicalxpress.com/news/2011-02-effective-treatment-lyme-disease-related-arthritis-proper.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.