

Heavy drinking in older teenagers has long- and short-term consequences

February 8 2011

In a systematic review of current evidence published in this week's *PLoS Medicine*, the authors—Jim McCambridge from the London School of Hygiene & Tropical Medicine, London, UK, and colleagues—conclude that there is enough evidence to recommend that reducing drinking during late adolescence is likely to be important for preventing long-term adverse consequences of drinking, as well as protecting against more immediate harms.

Although there is an urgent need for better studies in this area, research to date provides some evidence that high [alcohol consumption](#) in late adolescence often continues into adulthood and is associated with long-term alcohol problems, including dependence.

The authors of the current study conducted a comprehensive literature review to identify 54 relevant studies which included at least one quantitative measure of the effects of alcohol, on outcomes in adulthood such as death, alcohol dependence, criminal offences, mental health, educational attainment, and smoking. The majority of these studies were multiple reports from ten cohorts, half of which were from the US.

The authors found that although there is consistent evidence that higher alcohol consumption in late adolescence continues into adulthood and is associated with alcohol and other problems, most of these studies could not strongly support direct causality because of their weak designs. Furthermore, although a number of studies suggested links with late adolescent drinking to adult physical and mental health and social

consequences, this evidence is generally of poorer quality and insufficient to infer causality.

According to the authors: "It is clear that the evidence base on long-term consequences is not as extensive nor as compelling as it could be." Despite this limitation, they are able to say: "late adolescent alcohol consumption appears a probable cause of increased drinking well into adulthood, through to ages at which adult social roles have been achieved." However, they caution: "Heavier [drinking](#) seems most likely, however, to be only one component in a complex causal process, whose contribution has probably been overestimated in previous studies because of uncontrolled confounding, setting aside the uncertainties induced by self-reported data."

More information: McCambridge J, McAlaney J, Rowe R (2011) Adult Consequences of Late Adolescent Alcohol Consumption: A Systematic Review of Cohort Studies. *PLoS Med* 8(2): e1000413. [doi:10.1371/journal.pmed.1000413](https://doi.org/10.1371/journal.pmed.1000413)

Provided by Public Library of Science

Citation: Heavy drinking in older teenagers has long- and short-term consequences (2011, February 8) retrieved 20 April 2024 from <https://medicalxpress.com/news/2011-02-heavy-older-teenagers-long-short-term.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--