

Hyperactive nerve cells may contribute to depression

February 23 2011

(PhysOrg.com) -- Scientists at the U.S. Department of Energy's (DOE) Brookhaven National Laboratory, Cold Spring Harbor Laboratory, and the University of California, San Diego School of Medicine, have identified hyperactive cells in a tiny brain structure that may play an important role in depression. The study, conducted in rats and appearing in the February 24, 2011, issue of *Nature*, is helping to reveal a cellular mechanism for depressive disorders that could lead to new, effective treatments.

The research provides evidence that inhibition of this particular brain region - the lateral habenula - using implanted electrodes can reverse certain behaviors associated with depression, and also provides a mechanism to explain this effect. These findings lend support to the use of deep brain stimulation as a clinical treatment for people with long-standing, treatment-resistant depression.

"This research identifies a new anatomical circuit in the brain that mediates depression, and shows how it interacts with the brain's reward system to trigger a constant disappointment signal - which certainly would be depressing," said Fritz Henn, a neurobiologist and psychiatrist at Brookhaven and Cold Spring Harbor laboratories and a co-investigator on the research. "But," he added, optimistically, "identifying this circuit and how it works may open new doors to reversing these effects."

For example, said co-investigator Roberto Malinow, a professor of neurosciences at the UCSD School of Medicine, "it's possible that the

genes specifically expressed in these neurons could be targeted genetically or pharmacologically in order to manipulate them and reduce depression."

Scientists have known that cells in the lateral habenula are activated by negative or unpleasant events, including punishment and disappointment, such as when you don't get an expected reward. It may seem intuitive that such negative stimuli can lead to depression, but not everyone who experiences disappointment collapses into a state of helplessness. To explore this connection, the scientists wanted to take a closer look at the brain circuits.

They examined the sensitivity of lateral habenula brain cells - particularly those that connect and send signals to the brain's reward centers - in two animal models of "learned helplessness," a form of depression, as well as in control animals that weren't helpless.

Overall, the scientists found that these lateral habenula nerve cells were hyperactive in the depressed animals but not in the controls. Furthermore, the degree of hyperactivity coincided with the degree of helplessness.

"The activation of the lateral habenula is known to influence the release of serotonin and norepinephrine, two targets of current antidepressant medications," said Henn. "The current study looked at the role of the lateral habenula in terms of the dopamine system, the system involved in reward signaling. We found that hyperactivity in the lateral habenula due to stress-induced helplessness shuts off the brain's reward system."

To explore whether electrical stimulation could potentially reverse this reward-dampening effect, the researchers placed a stimulating electrode in the lateral habenula and measured the effects on the brain cells leading to the reward center. This was a study of rat brain cells that

simulated the effects of deep brain stimulation, a technique that is currently being explored as a treatment for clinical depression, which has shown promising results. The scientists found that electrical stimulation of hyperactive habenula [brain cells](#) markedly decreased excitatory activity leading to the reward center.

Next the scientists tested to see if deep brain stimulation in living rats that exhibited helplessness would affect their behavior. The result was a marked reduction in helpless behavior that was dependent on both placement of the electrode in the lateral habenula (not adjacent brain regions), and the intensity of the stimulation.

"Our results clearly show that suppression of synaptic transmission at the lateral habenula through deep brain stimulation can acutely reverse helpless behavior in rats," said Henn. "It's very likely that this beneficial effect was mediated by a suppression of excitatory nerve cells leading to the brain's reward system, as we observed in the cellular studies."

"Our study provides a cellular mechanism that may explain the hyperactivity of lateral habenula [nerve cells](#) observed in depressed humans and animal models of depression, as well as why 'silencing' these circuits, whether surgically or pharmacologically, can reduce depression-like symptoms in animals," Henn said.

Identifying these specific brain circuits and their dysfunction in depression may open the door to new effective treatments, including, potentially, lateral-habenula-targeted [deep brain stimulation](#).

More information: Synaptic potentiation onto habenula neurons in the learned helplessness model of depression:

www.nature.com/nature/journal/...ull/nature09742.html

Provided by Brookhaven National Laboratory

Citation: Hyperactive nerve cells may contribute to depression (2011, February 23) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2011-02-hyperactive-nerve-cells-contribute-depression.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.