

Researchers find reduced levels of an important neurotransmitter in MS

February 10 2011

(PhysOrg.com) -- Researchers at the University of Illinois at Chicago have shown for the first time that damage to a particular area of the brain and a consequent reduction in noradrenaline are associated with multiple sclerosis.

The study is available online in the journal [Brain](#).

The pathological processes in MS are not well understood, but an important contributor to its progression is the infiltration of [white blood cells](#) involved in immune defense through the blood-brain barrier.

Douglas Feinstein, research professor in [anesthesiology](#) at the UIC College of Medicine, and his colleagues previously showed that the neurotransmitter noradrenaline plays an important role as an immunosuppressant in the brain, preventing [inflammation](#) and stress to neurons. Noradrenaline is also known to help to preserve the integrity of the blood-brain barrier.

Because the major source of noradrenaline is neurons in an area of the brain called the locus coeruleus, the UIC researchers hypothesized that damage to the LC was responsible for lowered levels of noradrenaline in the brains of MS patients.

"There's a lot of evidence of damage to the LC in Alzheimer's and Parkinson's disease, but this is the first time that it has been demonstrated that there is stress involved to the neurons in the LC of MS

patients, and that there is a reduction in brain noradrenaline levels," said Paul Polak, research specialist in the health sciences in anesthesiology and first author on the paper.

For the last 15 years, Feinstein and his colleagues have been studying the importance of noradrenaline to inflammatory processes in the brain.

"We have all the models for studying this problem, so in some ways it was a small step to look at this question in MS," said Polak.

The researchers found that LC damage and reduced levels of noradrenaline occur in a mouse model of MS and that similar changes could be found in the brains of MS patients.

The findings suggest that LC damage, accompanied by reduction in noradrenaline levels in the brain, may be a common feature of neurologic diseases, Polak said.

"There are a number of FDA-approved drugs that have been shown to raise levels of noradrenaline in the brain, and we believe that this type of therapeutic intervention could benefit patients with MS and other neurodegenerative diseases, and should be investigated," he said.

Provided by University of Illinois at Chicago

Citation: Researchers find reduced levels of an important neurotransmitter in MS (2011, February 10) retrieved 9 April 2024 from <https://medicalxpress.com/news/2011-02-important-neurotransmitter-ms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
