

# Researchers reveal first autism candidate gene that demonstrates sensitivity to sex hormones

February 16 2011

---

George Washington University researcher, Dr. Valerie Hu, Professor of Biochemistry and Molecular Biology, and her team at the School of Medicine and Health Sciences, have found that male and female sex hormones regulate expression of an important gene in neuronal cell culture through a mechanism that could explain not only higher levels of testosterone observed in some individuals with autism, but also why males have a higher incidence of autism than females.

The gene, RORA, encodes a protein that works as a "master switch" for [gene expression](#), and is critical in the development of the cerebellum as well as in many other processes that are impaired in autism. Dr. Hu's earlier research found that RORA was decreased in the autistic brain. In this study, the research group demonstrates that aromatase, a protein which is regulated by RORA, is also reduced in autistic brains.

This is significant because aromatase converts testosterone to estrogen. Thus, a decrease in aromatase is expected to lead in part to build up of [male hormones](#) which, in turn, further decrease RORA expression, as demonstrated in this study using a neuronal cell model. On the other hand, female hormones were found to increase RORA in the [neuronal cells](#). The researchers believe that females may be more protected against RORA deficiency not only because of the positive effect of estrogen on RORA expression, but also because estrogen receptors, which regulate some of the same [genes](#) as RORA, can help make up for

the deficiency in RORA.

"It is well known that males have a higher tendency for autism than females; however, this new research may, for the first time, provide a molecular explanation for why and how this happens. This is just the tip of the iceberg in terms of understanding some of the biology underlying autism, and we will continue our work to discover new ways to understand and, hopefully, to someday combat this [neurodevelopmental disorder](#)," said Dr. Hu.

In her research published in 2009, Dr. Hu and colleagues found that RORA deficiencies were only apparent in the most severe cases of [autism](#) and were observed in the brain tissues of both male and female subjects. They further found that the deficiency in RORA was linked to a chemical modification of the gene (called methylation) which effectively reduces the level of RORA.

Provided by George Washington University Medical Center

Citation: Researchers reveal first autism candidate gene that demonstrates sensitivity to sex hormones (2011, February 16) retrieved 20 March 2024 from <https://medicalxpress.com/news/2011-02-reveal-autism-candidate-gene-sensitivity.html>

|   |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|