

Reducing the toll of alcohol in Canada

February 7 2011

Focused programs and public health policies can help reduce the burden of alcohol in Canada, which contributes significantly to acute and chronic diseases, social problems and trauma, states an analysis in *CMAJ* (*Canadian Medical Association Journal*).

The analysis, by researchers at the Centre for Addiction and Mental Health, Toronto, the Centre for Addictions Research of British Columbia and the University of Victoria, outlines several steps that must be taken to reduce the impact of alcohol in society and on individuals.

The estimated direct [health care costs](#) of alcohol in Canada in 2002 were \$3.3 billion and the total direct and indirect costs were \$14.6 billion compared with \$17 billion from tobacco and \$8.2 from illegal drugs. Per capita sales have risen 13% since 1996 and are expected to continue rising. In 2002, an estimated 450 000 Canadians were dependent drinkers, 1.3 million were high-risk drinkers and there were more than 8300 alcohol-related deaths.

To reduce the toll of alcohol, the authors recommend a comprehensive public health approach with population-level policies, including, for example, prices that are based on alcohol strength and keep pace with inflation, eliminating discount pricing, strengthening government control systems, restricting geographic access, increasing effectiveness of drunk-driving penalties and encouraging health practitioners to deliver brief interventions. They also call for limits on alcohol marketing and promotion.

"An effective response must be multidimensional, involving a combination of population-level policies, targeted interventions and special services for those who are high-risk drinkers or dependent on alcohol," writes Dr. Norman Giesbrecht, Centre for Addiction and Mental Health, Toronto, with coauthors.

The authors cite evidence that demonstrates the effectiveness of these interventions in other countries.

"Alcohol is often not 'on the radar screen' as a major public health issue in the broader public health community," they write. They argue that coordinated, broad-based action is required from physicians, [public health](#) officials, disease prevention alliances, charities, medical associations and government to help raise awareness of this issue and to help reduce the health and safety effects of [alcohol](#) in Canada.

Provided by Canadian Medical Association Journal

Citation: Reducing the toll of alcohol in Canada (2011, February 7) retrieved 18 April 2024 from <https://medicalxpress.com/news/2011-02-toll-alcohol-canada.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--