


In adolescence, the power to resist blooms in the brain

March 9 2011



(PhysOrg.com) -- Just when children are faced with intensifying peer pressure to misbehave, regions of the brain are actually blossoming in a way that heighten the ability to resist risky behavior, report researchers at three West Coast institutions.

The findings -- detailed in the March 10 issue of the journal *Neuron* -- may give parents a sigh of relief regarding their kids as they enter adolescence and pay more attention to their friends. However, the research provides scientists with basic insight about the brain's wiring,

rather than direct clinical relevance for now.

In the study, 24 girls and 14 boys from ethnically and socioeconomically diverse backgrounds underwent [functional magnetic resonance imaging](#) (fMRI) scans twice, at ages 10 and 13, the latter representing when children have moved into early adolescence. Each time, they were presented with photos of faces making neutral, angry, fearful, happy and sad emotional expressions.

Non-invasive fMRI, when focused on the brain, measures blood flow changes using a magnetic field and radio frequency pulses, producing detailed images that provide scientists with information about [brain activity](#) or help medical staff diagnose disease.

Researchers compared the fMRI results from age 10 to age 13, finding that activity increased significantly in the ventral striatum and the ventral medial portion of the [prefrontal cortex](#) over this three-year period. In addition to the scans, the researchers considered the children's self-reports on their ability to resist peer influences and engagement in risky or delinquent behavior.

The most enhanced response occurred in the ventral striatum, a brain region most frequently associated with reward-related processing. Over time, increases in brain activity there correlated with increases in children's resistance to peer influence.

"This is a complex point, because people tend to think of adolescence as the time when teenagers are really susceptible to [peer pressure](#)," said Jennifer H. Pfeifer, professor of psychology at the University of Oregon. "That is the case, but in addition to that added susceptibility they are also improving their ability to resist it. It's just that peer pressure is increasing because they spend a lot more time with peers during this time and less time with family. So it is a good thing that resistance to such influences

is actually strengthening in their brains."

This study, which researchers believed to be the first to report longitudinal fMRI findings about changes in the way the brain processes emotion during this critical time of brain development, appears to fit into a growing body of evidence that ventral striatum development during early adolescence is critical to emotional regulation carried out by the brain's prefrontal circuitry, the researchers concluded.

"This is basic research that hopefully is laying the foundation for future studies with even more clinical relevance," said Pfeifer, director of the Developmental Social Neuroscience Lab. "We really have a lot to learn about how the brain responds to really basic emotional stimuli across development."

There was a surprise finding that deserves more study, though, Pfeifer said. Responses in the amygdala -- a small almond-shaped mass centrally located deep in the brain -- showed significant increases during this period only to the sad faces.

The amygdala plays a major role in emotional reactivity and indexing the salience of things in the environment. It's possible, Pfeifer said, that this response to sad faces could somehow be tied to the emergence of depression, especially in girls.

"This response in the amygdala raises questions we hope to pursue," she said. "The span from age 9 to 13 is critical in pubertal development. How do individual differences apply here? Identifying this response to 'sadness' in the amygdala opens the door to thinking about how changes in emotional reactivity might be related to the increase in depression that we see as kids enter puberty. Rates of depression are particularly enhanced for teen girls. Is this increased response to sad faces somehow part of that?"

Based on results of the new study, she added, "I think what we know about the ventral striatum may be poised to undergo a transformation over the next several years."

Provided by University of Oregon

Citation: In adolescence, the power to resist blooms in the brain (2011, March 9) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2011-03-adolescence-power-resist-blooms-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.