

# Depression may increase the risk of kidney failure

March 10 2011

---

Depression is associated with an increased risk of developing kidney failure in the future, according to a study appearing in an upcoming issue of the *Clinical Journal of the American Society Nephrology* (CJASN). Approximately 10% of the US population will suffer from depression at some point during their lifetime.

Lead investigator, Dr. Willem Kop (Department of Medical Psychology and Neuropsychology at the University of Tilburg, the Netherlands) and colleagues studied 5,785 people from four counties across the United States for 10 years. The participants were 65 years and older and not yet on dialysis. They completed a questionnaire measuring depressive symptoms and a broad range of medical measurements, including estimated glomerular filtration rate (eGFR) and risk factors for kidney and heart diseases. The investigators examined whether depression predicted the onset of kidney disease or other medical problems in which the kidneys play a critical role.

According to the results, depression coincided with the presence of chronic kidney disease (CKD) and was 20% more common in individuals with kidney disease than those without kidney disease. The study shows that depression predicted subsequent rapid decline in kidney function, new onset clinically severe kidney disease (or end-stage renal disease), and hospitalizations that were complicated by acute kidney injury. When the investigators corrected for the long-term effects of other medical measures, the predictive value of depression for hospitalizations with acute kidney injury remained high.

Take home message: "People with elevated [depressive symptoms](#) have a higher risk of subsequent adverse kidney disease outcomes. This is partially explained by other medical factors related to depression and [kidney disease](#). But, the association with depression was stronger in patients who were otherwise healthy compared to those who had co-existing medical disorders such as diabetes or heart disease," explains Kop.

The investigators are currently analyzing which factors may explain the association with depression, which could include delayed seeking of medical care and miscommunications between patient and physicians and important biological processes associated with [depression](#), such as the immune and nervous systems.

**More information:** The article, entitled "Longitudinal Association of Depressive Symptoms with Rapid Kidney Function Decline and Adverse Clinical Renal Disease Outcomes," will appear online at [doi:10.2215/CJN.03840510](https://doi.org/10.2215/CJN.03840510)

Provided by American Society of Nephrology

Citation: Depression may increase the risk of kidney failure (2011, March 10) retrieved 24 April 2024 from <https://medicalxpress.com/news/2011-03-depression-kidney-failure.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--