

Doctors lax in monitoring potentially addicting drugs

March 3 2011

Few primary care physicians pay adequate attention to patients taking prescription opioid drugs -- despite the potential for abuse, addiction and overdose, according to a new study by researchers at Albert Einstein College of Medicine of Yeshiva University.

The study, published in the March 2 online edition of the [Journal of General Internal Medicine](#), found lax monitoring even of patients at high risk for opioid misuse, such as those with a history of [drug abuse](#) or dependence. The findings are especially concerning considering that prescription drug abuse now ranks second (after marijuana) among illicitly used drugs, with approximately 2.2 million Americans using pain relievers nonmedically for the first time in 2009, according to the National Institute on Drug Abuse (NIDA).

"Our study highlights a missed opportunity for identifying and reducing misuse of prescribed [opioids](#) in primary care settings," said lead author Joanna Starrels, M.D., M.S., assistant professor of medicine at Einstein. "The finding that physicians did not increase precautions for patients at highest risk for opioid misuse should be a call for a standardized approach to monitoring."

The researchers studied administrative and medical records of more than 1,600 primary care patients for an average of two years while they received regular prescription opioids for chronic, non-cancer pain. They looked at whether patients received urine drug testing, were seen regularly in the office, or received multiple early opioid refills.

Only a small minority (8 percent) of patients were found to have undergone any urine drug testing. While such testing was more common in patients at higher risk for opioid misuse, the rate of testing among those high-risk patients was still low (24 percent). Only half of patients were seen regularly in the office, and patients at higher risk of opioid misuse were not seen more frequently than patients at lower risk. Although fewer than one-quarter (23 percent) of all patients received two or more early opioid refills, patients at greater risk for opioid misuse were more likely to receive multiple early refills.

"We were disturbed to find that patients with a drug use disorder were seen less frequently in the office and were prescribed more early refills than patients without these disorders," said Dr. Starrels. "We hope that these findings will call attention to this important safety concern."

Prescription drug misuse is a major public health problem. In a 2004 NIDA [report](#), it was estimated that 48 million people over the age of 12 have taken prescription drugs for nonmedical uses in their lifetime – which represents approximately 20 percent of the U.S. population. Opioids, central nervous system depressants and stimulants were the drugs most commonly abused.

"Most [primary care physicians](#) are attuned to these problems," said Dr. Starrels, "but they haven't put sufficient strategies in place to help reduce risks." She and her co-authors recommend that physicians adopt the following risk-reduction strategies: standardize a plan of care for all patients on long-term opioids, which includes urine drug testing; schedule regular face-to-face office visits to evaluate patients' response to opioids and evidence of misuse; and stick to a previously agreed-upon refill schedule.

Provided by Albert Einstein College of Medicine

Citation: Doctors lax in monitoring potentially addicting drugs (2011, March 3) retrieved 7 May 2024 from <https://medicalxpress.com/news/2011-03-doctors-lax-potentially-addicting-drugs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.