

## Stroke incidence higher among patients with certain type of retinal vascular disease

March 14 2011

---

Patients with a disease known as retinal vein occlusion (RVO) have a significantly higher incidence of stroke when compared with persons who do not have RVO, according to a report in the March issue of *Archives of Ophthalmology*, one of the JAMA/Archives journals.

"[Retinal vein occlusion](#) (RVO) is a retinal vascular disease in which a retinal vein is compressed by an adjacent retinal artery, resulting in [blood flow](#) turbulence, thrombus formation, and retinal ischemia," the authors write as background information in the article. "Although RVO is a significant cause of severe visual impairment in adults, it can occur at any age." Older age, diabetes, hypertension and vascular disease are among the [risk factors](#) for RVO.

Winifred Werther, Ph.D., then of Genentech Inc., South San Francisco, Calif., now of Vertex Pharmaceuticals, Cambridge, Mass., and colleagues conducted a retrospective cohort study to compare the incidence rates of [myocardial infarction](#) (MI, [heart attack](#)) and cerebrovascular accident (CVA, stroke) in hospitalized patients with and without retinal vein occlusion. The researchers used a U.S. population-based health care claims database to identify patients with RVO and control patients, matched for age and sex.

Among 4,500 patients with RVO and 13,500 control patients, the researchers found that patients with RVO had an almost two-fold higher incidence of stroke than the age- and sex-matched controls.

"Event rates for CVA were 1.16 and 0.52 per 100 person-years for RVO and controls, respectively," the authors report.

In contrast, event rates for heart attack were similar in the patients with RVO and the control patients.

"Although men and patients younger than 65 years with RVO had a 1.6- and 1.9-fold higher risk of MI, respectively, compared with controls, there were no statistically significant differences in MI rates between patients with RVO and controls when they were stratified by sex or age," the authors write.

They conclude that "these data suggest that physicians and patients should be aware of the possible increased risk of CVA but not of MI in patients with RVO."

**More information:** Arch Ophthalmol, 2011;129[3]:326-331

Provided by JAMA and Archives Journals

Citation: Stroke incidence higher among patients with certain type of retinal vascular disease (2011, March 14) retrieved 26 April 2024 from <https://medicalxpress.com/news/2011-03-incidence-higher-patients-retinal-vascular.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.