

Laughter really is the best medicine (for leg ulcers)

March 8 2011

Forget technology. The best prescription for patients with venous leg ulcers is good quality nursing care – and the occasional belly laugh!

A five-year study led from the University of Leeds has shown that ultrasound therapy does nothing to speed up the healing process of leg ulceration – contrary to what had been expected. Traditional methods of nursing care, which are cheaper and easier to deliver, work just as well, the authors conclude.

"The 'healing energy' of low-dose ultrasound can make a difference to some medical conditions but with venous leg ulcers, this is simply not the case," said Professor Andrea Nelson from the University of Leeds' School of Healthcare, who led the study.

"The key to care with this group of patients is to stimulate blood flow back up the legs to the heart. The best way to do that is with compression bandages and support stockings – not 'magic wands' - coupled with advice on diet and exercise. Believe it or not, having a really hearty chuckle can help too. This is because laughing gets the diaphragm moving and this plays a vital part in moving blood around the body."

Venous leg ulcers are common in people with varicose veins or mobility problems whose 'muscle pumps' in the feet and calves struggle to drive blood up to the heart. These ulcers can be painful and unsightly, having a significant negative impact on health and quality of life.

Although most leg ulcers will clear up with good nursing care, a significant proportion of the lesions will take 12 months or longer to heal. The older and larger ulcers become the harder they are to get rid of, hence the search for solutions that could speed-up the healing process.

A number of small studies had suggested that ultrasound could be the answer. Professor Nelson (University of Leeds), working with colleagues from the Universities of York and Teeside, and many NHS Trusts, have now shown that this is not the case.

The researchers concentrated on 'hard to heal' ulcers that had not cleared up after six months or longer. Drawing on patients from across the UK and Ireland, they found that adding ultrasound to the standard approach to care - dressings and compression therapy - made no difference to the speed of healing or the chance of the ulcers coming back. Ultrasound also raised the cost of care per patient by almost £200.

"Rising levels of obesity mean that the number of people who suffer from legs ulcers is likely to grow," Professor Nelson said. "We do need to find ways to helping those patients who ulcers won't go away, but our study shows that ultrasound is not the way to do that. We need to focus on what really matters, which is good quality nursing care. There really is no need for the NHS to provide district nurses with ultrasound machines. This would not be money well spent."

More information: 'VenUS III: a randomised controlled trial of therapeutic ultrasound in the management of venous leg ulcers' is available online in advance of publication in Health Technology Assessment [DOI:10.3310/hta15130](https://doi.org/10.3310/hta15130)

Provided by University of Leeds

Citation: Laughter really is the best medicine (for leg ulcers) (2011, March 8) retrieved 3 May 2024 from <https://medicalxpress.com/news/2011-03-laughter-medicine-leg-ulcers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.