

New national study finds boxing injuries on the rise; youth head injury rates also concerning

March 2 2011

The risk and nature of injury in the sport of boxing has generated a great deal of controversy in the medical community, especially in relation to youth boxing. A new study, conducted by researchers in the Center for Injury Research and Policy of The Research Institute at Nationwide Children's Hospital, examined boxing injuries among participants 6 years of age and older from 1990 to 2008.

During the 19-year study period, an average of 8,700 [boxing](#) injuries were treated in United States emergency departments each year, and approximately 2,500 of those injuries were to children and [adolescents](#) 6 to 17 years of age. The number of boxing injuries each year increased 211 percent during the study, climbing from 5,361 injuries in 1990 to nearly 17,000 injuries in 2008.

According to the study, released online February 28, 2011 by the [American Journal of Preventive Medicine](#), the most common injury diagnosis was a fracture (28 percent). The hand was the most frequently injured body region (33 percent), followed by the head and neck (23 percent). While the majority of injuries occurred at a sports or recreation facility (54 percent), one-third of the injuries (34 percent) occurred at home.

The most concerning discovery from the study was the similar proportion of concussions/closed head injuries (CHIs) among the age

groups (9 percent among 12-17 year olds, 8 percent among 18-24 year olds and 9 percent among 25-34 year olds).

"We expected a smaller proportion of concussions/CHIs among younger boxers, since they generate a lower punch force," said Gary Smith, MD, DrPH, senior author of the study and director of the Center for Injury Research and Policy at Nationwide Children's Hospital. "The fact that young boxers are experiencing a similar proportion of concussions and CHI's as older boxers is extremely concerning given the potential risk of developing chronic traumatic encephalopathy (CTE) with repetitive brain trauma. These repetitive blows to the head may be placing boxers under 18 years of age at risk for neurological impairment and psychological problems due to CTE."

The findings from this study support the position of medical societies that oppose boxing, especially among youth.

"Although there is risk of injury with most sports, boxing is unique because participants are rewarded for intentionally striking their opponent in the face and head with the intent of harming or incapacitating them," said Dr. Smith, also a professor of Pediatrics at The Ohio State University College of Medicine. "The increasing number of boxing injuries, coupled with the potential long-term consequences of these injuries, suggests that increased injury prevention efforts are needed."

This is the first nationally representative study to examine boxing injuries treated in emergency departments. Data for this study were obtained from the National Electronic [Injury](#) Surveillance System (NEISS), which is operated by the U.S. Consumer Product Safety Commission. The NEISS dataset provides information on consumer product-related and sports and recreation-related injuries treated in hospital emergency departments across the country. Boxing participation

data were obtained from the Superstudy of Sports Participation.

Provided by Nationwide Children's Hospital

Citation: New national study finds boxing injuries on the rise; youth head injury rates also concerning (2011, March 2) retrieved 13 March 2024 from

<https://medicalxpress.com/news/2011-03-national-injuries-youth-injury.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|