

# For back, neck pain, artificial disc replacement has cost, outcome advantages over fusion surgery

March 22 2011

---

When physical therapy and drugs fail to relieve back or neck pain, patients often turn to spinal fusion surgery as a last resort, but two new studies show that in certain situations, especially when several discs are involved, artificial disc replacement may give better long-term results at lower cost.

Hospital costs for artificial disc replacement were 49 percent lower, and four years out from surgery, artificial disc patients were four times less likely to need additional surgery than those who had fusion operations, according to two recently published studies.

"Back pain is the fifth leading cause of [hospital admission](#) and the third most common reason for surgery," said orthopaedic surgeon Rick B. Delamarter, M.D., co-director of the Cedars-Sinai Spine Center.

"Estimates vary and are probably understated, but health care expenditures for back pain top \$91 billion a year, not including indirect and societal costs such as time lost from work and worker's compensation. It is crucial that we develop surgical procedures that are cost effective without sacrificing high-quality results."

Delamarter is a lead author of the two studies, which were published recently in the SAS Journal of the International Society for the Advancement of Spine Surgery and the Society for Minimally Invasive Spine Surgery.

Both studies compared disc replacement surgery with the more common fusion operation to treat degenerative disc disease – deterioration caused by aging and wear and tear. One study looked at 209 patients with damaged neck discs who received either minimally invasive disc replacement or the more complex [spinal fusion](#) surgery. These patients were followed at regular intervals for four years. A separate group of 136 who received an artificial disc two years after the first group also were part of the study. All patients were assessed on their satisfaction with the results of the procedure. The other study focused on 53 patients suffering from three-level, lower back disc disease and looked at cost comparisons for length of hospital stay, resources used and other factors.

Discs act as cushions between the bones (vertebrae) of the spine. When healthy, the discs have enough "give" to allow the back to be flexible but are firm enough to provide stability. With age or injury, they lose their pliability and density. Nerves may become pinched between the bones, causing pain not just in the spine but in other parts of the body.

Fusion surgery seeks to relieve symptoms of the degenerative disease by removing the damaged disc and replacing it with bone. Studies show this procedure often can be effective but there can be drawbacks: in some cases, fused spinal sections can lose flexibility, impeding normal movement and stressing adjacent discs, often leading later to more fusion surgery. Artificial disc replacement, which has been performed in the United States since 2000, tends to cause less tissue injury than fusion surgery, and the discs are designed to maintain natural spine movement and reduce need for follow-up surgery.

In their study of patients suffering neck (cervical) disc disease, Delamarter and his colleagues found both disc replacement and fusion surgery acceptable options providing good long-term outcomes. But four years out from surgery, fusion patients were four times more likely to need more surgery. Half of these secondary operations were needed to

treat new disc problems occurring adjacent to fused sections.

Patients in the study had imaging scans to measure before-and-after flexibility and range of motion; were evaluated for physical and neurological improvement; and completed self-assessment questionnaires on neck disability, neck and arm pain intensity. Improvements were seen in all categories, regardless of the [surgical procedure](#), but results tended to be at least slightly better in patients with disc replacements.

An important area, according to Delamarter, was patient satisfaction with the procedure. Four years after surgery, nearly 88 percent of disc replacement patients reported that they were very to completely satisfied, compared with 76 percent of fusion patients. When asked if they would undergo the same surgery again, 89 percent of disc replacement patients and 81 percent of fusion patients said yes.

"While the results of this 13-center study are preliminary, it appears that artificial disc replacement surgery compares favorably as an effective procedure, is preferred by patients and slows the rate of adjacent-level disease," Delamarter said.

Spinal fusion has been recognized as one of the more expensive surgical procedures. The second study looked at costs and is believed to be the first to compare those of three-level disc replacement with three-level fusion. It evaluated itemized billing records of 53 patients undergoing surgery for three consecutive discs of the low back (lumbar spine) at St. John's Health Center in Santa Monica, Calif., where Delamarter practiced before joining Cedars-Sinai.

Total hospital costs for patients undergoing disc replacement surgery averaged 49 percent lower than those for fusion patients

The number of fusions for low back pain is rising rapidly, but Delamarter urges hospitals, insurers and surgeons to consider alternatives, including total disc replacement.

"After safety and effectiveness of a procedure have been documented, cost is an important consideration," he said. "Artificial disc replacement technology appears to offer a promising, cost-saving alternative to fusion, especially for patients with disease at three levels."

**More information:** Citation: SAS Journal, December 2010 "Results at 24 months from the prospective, randomized, multi-center Investigational Device Exemption trial of ProDisc-C versus anterior cervical discectomy and fusion with 4-year follow-up and continued access patients," and "Cost comparison of patients with 3-level artificial total lumbar disc replacements versus 360-degree fusion at 3 contiguous lumbar vertebral levels: an analysis of compassionate use at 1 site of the US investigational device exemption clinical trial."

Provided by Cedars-Sinai Medical Center

Citation: For back, neck pain, artificial disc replacement has cost, outcome advantages over fusion surgery (2011, March 22) retrieved 26 April 2024 from <https://medicalxpress.com/news/2011-03-neck-pain-artificial-disc-outcome.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.