

More tests needed for oesophageal cancer patients

March 1 2011

A University of Adelaide medical researcher says current treatment for people diagnosed with oesophageal cancer could be improved with additional pathology tests.

Dr Sarah Thompson, a surgeon and PhD candidate, says tiny cancer cells are being missed in routine pathology examinations, resulting in half of patients who have been given the "all-clear" dying within five years.

In a study of 250 oesophageal cancer patients in Adelaide over the past 10 years, Dr Thompson found that of the 119 people given a clean bill of health, only 62 were still alive five years later.

"If you tell people they are totally clear of cancer and then a year later they come back with secondary cancer, that is a devastating result," she says.

More than 1000 people are diagnosed with cancer of the oesophagus in Australia each year. There are two types: [squamous cell carcinoma](#), occurring in the top part of the oesophagus; and [adenocarcinoma](#), which is closer to the stomach. The latter has increased by 400% in Australia since the 1970s.

Dr Thompson's study showed that oesophageal tumour cells are being missed in the [lymph nodes](#) in conventional pathological tests.

"Pathologists normally take just one cut from a lymph node when testing

for cancer and this is usually sufficient. But our research has shown that in one third of cancer patients with negative nodes, small cancer cells will be present. These are only visible with special stains and at least three cuts of the node," Dr Thompson says.

"A controversy in the medical world, as far as predicting cancer survival rates, is whether or not those tiny, single [cancer cells](#) mean anything. Some medical experts believe these cells are just in transit and passing through our [blood stream](#). Others think if they are present in the lymph nodes these patients should be offered chemotherapy as a matter of course."

Dr Thompson says the oesophageal cancer patients who have the most to gain are those who have had surgery without chemotherapy or radiation beforehand to shrink the tumour.

"Chemotherapy after surgery might improve their survival."

"I think there is also merit in changing the existing pathology guidelines for oesophageal [cancer patients](#) to ensure a more accurate diagnosis. Further sectioning and staining of the lymph nodes should be carried out if the lymph nodes are negative after only one cut to rule out the presence of isolated [tumour cells](#)," Dr Thompson says.

Her findings have been published in the [Annals of Surgery](#) in an article co-authored by Dr Thompson's PhD supervisor Professor Glyn Jamieson, the Dorothy Mortlock Professor of Surgery at the University of Adelaide.

Provided by University of Adelaide

Citation: More tests needed for oesophageal cancer patients (2011, March 1) retrieved 19 April

2024 from <https://medicalxpress.com/news/2011-03-oesophageal-cancer-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.