

# Side effects of prophylactic percutaneous endoscopic gastrostomy placement

March 15 2011

---

A research team from United States determined the rate of use and non-use of prophylactic percutaneous endoscopic gastrostomy (PEG) tubes among patients with head and neck cancer patients. They found that there is a high rate of unnecessary PEG placement when done prophylactically in patients with head and neck cancer.

Impairment of oral intake occurs in the majority of patients with [head and neck cancer](#) (HNC) receiving chemoradiotherapy. Placement of prophylactic percutaneous endoscopic gastrostomy (PEG) tube in asymptomatic newly diagnosed HNC before chemoradiation is a common practice in some centers. In some studies, PEG has been associated with a decrease in treatment related weight loss in patients with HNC, but no studies have examined the utilization rate.

A research article published on February 28, 2011 in the [World Journal of Gastroenterology](#) addresses this question. The authors have observed anecdotally that a number of HNC patients who received a prophylactic PEG tube in fact never used them. They performed a retrospective database study of all patients in whom PEG tube was placed for HNC to determine the prevalence of unused prophylactically placed PEG tubes. Data were also analyzed for possible factors predictive of unused PEGs or PEGs used for less than 2 wk.

This is the first study that addressed the issue of use of prophylactic PEG in HNC patients. The result of this study showed that a significant number of patients (47%) with prophylactic PEG tubes never used their

PEG or used it for less than 2 wk. No association with PEG use vs non-use was observed for cancer diagnosis, stage, or specific [cancer treatment](#).

The study suggests prophylactic PEG placement prior to HNC therapy is associated with a high rate of non use or limited use. Further prospective studies evaluating specific selection criteria for prophylactic PEG in this setting are needed. Similarly, additional studies are needed to assess the impact of prophylactic PEG tube placement on the cost-effectiveness of cancer care, quality of life, hospital admission rate, and, most importantly, survival.

**More information:** Madhoun MF, Blankenship MM, Blankenship DM, Kreml GA, Tierney WM. Prophylactic PEG placement in head and neck cancer: How many feeding tubes are unused (and unnecessary)? World J Gastroenterol 2011; 17(8): 1004-1008.  
[www.wjgnet.com/1007-9327/full/v17/i8/1004.htm](http://www.wjgnet.com/1007-9327/full/v17/i8/1004.htm)

Provided by World Journal of Gastroenterology

Citation: Side effects of prophylactic percutaneous endoscopic gastrostomy placement (2011, March 15) retrieved 30 April 2024 from <https://medicalxpress.com/news/2011-03-side-effects-prophylactic-percutaneous-endoscopic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--