

Diabetes guidelines linked to severe low blood sugar in frail elderly

May 23 2011

When an independent senior health program implemented new recommended diabetes blood sugar guidelines, episodes of severe hypoglycemia (low blood sugar) tripled among frail elderly patients, according to a study led by Sei J. Lee, MD, a geriatrician at the San Francisco VA Medical Center.

The study, which was published in the April 2011 issue of the [Journal of the American Geriatrics Society](#), was the first to measure what happened when the guidelines, developed in 2003 by the American Geriatrics Society, were implemented among frail elders, said Lee. The study focused on outcomes at On Lok Senior Health, in San Francisco.

The guidelines call for a hemoglobin A1c level of less than 8 percent. The hemoglobin A1c test, which measures average [blood sugar](#) over the previous three months before the test, is considered an indicator of long-term [blood sugar control](#) for diabetics.

The long-term goal of such guidelines is the avoidance of vascular complications — including heart attack, stroke, and kidney disease — that are brought about by chronically high blood sugar, which damages circulation in small blood vessels. In order to accomplish this goal, blood sugar is monitored frequently during the day, and kept relatively low through diet, medications, and insulin injections — a regimen known as tight control.

When the frail elders were put on tight control, reported Lee, “there was

good news and bad news.” The good news, he said, was that the incidence of hyperglycemia — blood sugar higher than 400 — decreased dramatically compared to pre-guideline levels.

At the same time, episodes of severe [hypoglycemia](#) — blood sugars of less than 50 that required trips to the emergency room — increased threefold. “These episodes were quite dangerous for these seniors,” said Lee. “When blood sugar gets too low, patients can become confused, they can fall, they can even become comatose. It can become life-threatening very quickly.”

Significantly, the increase in hypoglycemic episodes occurred during the first 18 months of the guidelines’ implementation; for the two years after that, the rate was similar to what it was before the guidelines were introduced. One possible explanation, according to the authors, is that a small number of patients are especially at risk of severe hypoglycemia in response to even a modest tightening of control. Once those patients have a severe hypoglycemic episode and are identified, the rest are treated successfully with the aggressive regimen.

“This says that the period when you are first implementing tighter control is the time of greatest risk,” said Lee. “That is the time to require closer follow-up of patients in order to make sure that these adverse effects are not occurring.”

Lee noted that the study took place under “ideal conditions” for implementing tight control with a minimum of bad outcomes: “This is a program where patients are seen several times a week, and where health care providers know exactly what medications they are taking, what and how much they’re eating, and how much physical activity they’re getting.”

If the occurrence of severe hypoglycemia increased even under these

circumstances, said Lee, “then for the general population of the frail elderly, who are not monitored nearly as closely as our study population, the current guideline may be too aggressive.”

More information: [onlinelibrary.wiley.com/doi/10 ...
15.2011.03362.x/full](https://onlinelibrary.wiley.com/doi/10.1111/15.2011.03362.x/full)

Provided by University of California, San Francisco

Citation: Diabetes guidelines linked to severe low blood sugar in frail elderly (2011, May 23)
retrieved 24 May 2024 from <https://medicalxpress.com/news/2011-05-diabetes-guidelines-linked-severe-blood.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--