

Drug-resistance fears for deadly fungal disease

May 5 2011

Deadly human fungal infections caused by certain strains of *Aspergillus* fungi appear to be developing resistance to current drug treatments at an alarming rate, say scientists.

University of Manchester researchers, working with colleagues in Newark, USA, have developed a new test that can not only better diagnose *Aspergillus* infection, but can also spot signs of antifungal resistance to azoles – the class of drugs used to treat patients with aspergillosis.

Using the new test, which uses direct molecular detection rather than culturing the fungus in a Petri dish, the team found that 55% of aspergillosis patients had telltale signs known as 'markers' that indicated they had developed resistance to azoles. The findings compare to resistance rates of 28% carried out by the team just two years ago using traditional culturing methods.

Furthermore, the study – published in the prestigious US journal *Clinical Infectious Diseases* – discovered azole-resistance markers in threequarters of the small number of aspergillosis patients (eight) whom had never been treated with an azole, suggesting widespread dissemination of resistance.

"*Aspergillus* significantly worsens asthma symptoms and causes lifethreatening infections in those with long-term lung infections or damaged immune systems, such as chemotherapy and transplant patients



or people with HIV," said David Denning, Professor of Medicine and Medical Mycology at The University of Manchester and Director of the National Aspergillosis Centre at the University Hospital of South Manchester NHS Foundation Trust.

"Using an ultrasensitive, real-time test for *Aspergillus*, similar to the method used to diagnose HIV, MRSA and influenza, we have directly detected azole resistance in people with aspergillosis, without first culturing the fungus in a dish. The presence of *Aspergillus* was detected in many more samples than using traditional culture methods, and 55% were found to contain azole-resistance markers.

"This is an extraordinarily high rate of resistance, possibly related to fungicide use in agriculture – more than a third of 'pesticides' used by UK farmers are azoles – and long treatment courses in patients, so the findings have major implications for the sustainability of azoles for human antifungal therapy."

The azoles itraconazole (Johnson & Johnson), voriconazole (Pfizer) and posaconazole (Merck) have annual sales of more than \$1bn annually. Conventional diagnosis of aspergillosis is limited by poor culture yield, and so the true frequency of azole resistance has been unclear.

In this latest study, the researchers analyzed phlegm from patients with allergic and chronic lung disease caused by *Aspergillus* and found that almost twice the proportion of individuals tested had resistance markers in their sample compared to a Petri dish (or culture) study carried out by the team in 2008/9.

Professor Denning added: "Not only is molecular testing much more sensitive than conventional culture for diagnosis, but it enables testing for resistance, which until now has been impossible if cultures are negative. Given the rising frequency of resistance in *Aspergillus* in



northern Europe, China and the United States, this study provides key data for doctors to shift antifungal therapy in the face of <u>resistance</u>."

More information: Denning DW, Park S, Lass-Florl C, Fraczek MG, Kirwan M, Gore R, Smith J, Bueid A, Bowyer P, Perlin DS. High frequency triazole resistance found in non-culturable Aspergillus fumigatus from lungs of patients with chronic fungal disease. Clin Infect Dis 2011;52:1123-9.

2009 study: Bueid A, Howard SJ, Moore CB, Richardson MD, Harrison E, Bowyer P, Denning DW. Azole antifungal resistance in Aspergillus fumigatus - 2008 and 2009. J Antimicrob Chemother 2010;65:2116-8.

Provided by University of Manchester

Citation: Drug-resistance fears for deadly fungal disease (2011, May 5) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2011-05-drug-resistance-deadly-fungal-disease.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.