

# Effects of celiac disease on bone mineral density are pronounced in lumbar spine than femoral neck

May 26 2011

---

Patients with celiac disease are more than 4.5 times more likely to develop osteoporosis compared to healthy people in an age and gender matched cohort with no identifiable risk factors for osteoporosis, according to a study presented today at the EULAR 2011 Annual Congress.

The British study of 1,030 people compares the extent of [bone mineral density](#) (BMD)\* loss – a common measure of osteoporosis – in two physical locations in the body. The results demonstrated that the mean BMD of the lumbar spine in patients with coeliac disease was significantly different when compared to the control group (1.79 (95% CI 1.32, 2.44)), however not for the femoral neck vs. control (1.52 (95% CI 0.86, 2.67)).

"This is the first study that compares the extent of BMD loss at the lumbar spine and femoral neck in an age and sex matched cohort, and demonstrates that the detrimental effects of coeliac disease are most pronounced in the lower back compared to the hip joint," said Mr Oldroyd of Lancaster University School of Medicine. "These findings may be due to the fact that the bone in the lumbar spine is spongy, less dense and weaker in comparison to the femoral neck, causing it to be more susceptible to the detrimental effects of coeliac disease. There is greater research required to determine why this effect is seen and whether it can have future implications for treatment."

The authors from Lancaster and the Liverpool area collected results of DEXA scans (used to measure BMD) from 1,030 patients with coeliac disease between June 2004 and September 2007. Scan results of the lumbar spine and femoral neck of patients with coeliac disease were compared to a healthy, age and gender matched cohort, with no identifiable risk factors for [osteoporosis](#). Logistic regression analysis was used to investigate the association between coeliac disease and BMD results for the lumbar spine and femoral neck in the lowest tertile of the whole study cohort.

Provided by European League Against Rheumatism

Citation: Effects of celiac disease on bone mineral density are pronounced in lumbar spine than femoral neck (2011, May 26) retrieved 10 April 2024 from <https://medicalxpress.com/news/2011-05-effects-celiac-disease-bone-mineral.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------