

# Study shows the HPV vaccine Gardasil doesn't increase disease activity in SLE patients

May 26 2011

---

Results of a Chinese study showed that the HPV vaccine did not have significant effects on the number of disease flares or antibody measures in patients with inactive SLE receiving stable doses of medications after administration, and therefore was determined safe to use to prevent HPV in this group of patients. SLE, an autoimmune disorder, affects nine times as many women as men<sup>1</sup> and studies have shown that the rate of HPV in this group is significantly higher than in the healthy population.<sup>2</sup> Vaccination is therefore an important consideration in protecting SLE patients from HPV infection, which has been shown to be responsible for cervical cancer.

"Our study set out to investigate whether vaccination with Gardasil increased disease flares in patients with SLE," said Professor Chi Chiu Mok from the Tuen Mun Hospital, Hong Kong. "The causal relationship between vaccination and flare wasn't clear, however what we do know is that the rate of flares was not increased post vaccination, confirming that, in the cohort studied, Gardasil was safe for use."

In the duration of the six-month study, there were three mild/moderate mucocutaneous flares (flares that occur in mucous-lined areas of the body), which were all controlled with usual [treatment regimens](#). The rate of flare-ups observed in this study (rate: 0.08/patient/year) was numerically lower than the rate observed in a cohort of SLE patients observed over a 5 year period (0.10/patient/year) though the reason for

this is unknown.

Furthermore there were no significant changes in the levels of various antibody measures used to assess level of disease activity. Disease [flares](#) (measured by the Systemic Lupus Erythematosus National Assessment (SELENA) flare instrument), disease activity scores (measured by the [Systemic Lupus Erythematosus](#) Disease Activity Index (SLEDAI)) and physicians' global assessment (PGA) scores were also the same from baseline to two and six months.

Female patients who fulfilled four ACR criteria for SLE were recruited for the study. The 50 women were between the ages of 18 and 35 (mean age  $25.8 \pm 3.9$  years) and had received a stable dose of prednisolone and/or other immunosuppressives (the normal treatment for patients with SLE) within three months of the study. The Gardasil vaccine was given at baseline, two months and six months by intramuscular injection, and various disease scores and antibody measures were recorded and analysed.

#### **More information: References:**

1 PubMed Health. Systemic Lupus Erythematosus.  
[www.ncbi.nlm.nih.gov/pubmedhealth/PMH0001471/](http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0001471/) Accessed 10 May 2011

2 Brunk, D. SLE Patients Have Higher HPV Rates, Riskier Subtypes.  
Rheumatology News 2009;8(7):18

Provided by European League Against Rheumatism

Citation: Study shows the HPV vaccine Gardasil doesn't increase disease activity in SLE patients

(2011, May 26) retrieved 7 May 2024 from <https://medicalxpress.com/news/2011-05-hpv-vaccine-gardasil-doesnt-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.