

Hypertension control in Canada has improved significantly

May 16 2011

Treatment and awareness of hypertension in Canada have improved significantly in the last 25 years for community-dwelling adults, states an article in *CMAJ* (*Canadian Medical Association Journal*).

High blood pressure ([hypertension](#)) is a risk factor for vascular disease and mortality in developed countries but if managed properly, it is the most important modifiable risk factor. Recent studies in Canada have indicated improvements in the prescribing rates of drugs to treat hypertension and consequent decreases in [cardiovascular events](#) related to [high blood pressure](#).

A team of researchers looked at the measurement of blood pressure in people aged 20 to 79 not living in institutions who participated in either of two national health surveys. The analysis showed decreases in the number of hypertensive Canadians between 1992 and 2009 who were not being treated or not receiving adequate treatment to control their blood pressure. As well, the percentage of people who were unaware of their condition had dropped from 43% to 17%. Systolic [blood pressure levels](#) were lower in people with treated hypertension and in people without high blood pressure in 2009 compared to 1992.

"The management of hypertension among community-dwelling Canadian adults appears to have improved over the past two decades," writes Dr. Finlay McAlister, University of Alberta, with coauthors. "Notably, the rates of awareness, treatment and control documented in the ON-BP and the CHMS [surveys] are higher than those recently

reported from physical measures surveys done in the United States and elsewhere during the same periods."

People with high blood pressure and heart disease or cardiovascular or [cerebrovascular disease](#) were more likely to have their [blood pressure](#) under control. "This is encouraging, as it suggests that either Canadian physicians appropriately target more intensive therapy to patients at higher risk or that patients at higher risk are more compliant with their therapy," state the authors.

"However, despite marked improvements in rates for the control of hypertension over the past two decades, one-third of community-dwelling Canadian adults with hypertension still have blood pressures that are higher than the currently recommended targets, and cardiovascular disease remains the most common cause of premature death and disability in Canada," the authors conclude, stating we need to further improve management of hypertension.

Provided by Canadian Medical Association Journal

Citation: Hypertension control in Canada has improved significantly (2011, May 16) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2011-05-hypertension-canada-significantly.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--