

# Insomnia linked to high insulin resistance in diabetics

May 2 2011

---

In the largest study of its kind to establish a link between sleep and diabetes, researchers found that people with diabetes who sleep poorly have higher insulin resistance, and a harder time controlling the disease.

The findings, published in the June issue of *Diabetes Care*, suggest that poor sleep may contribute to worse outcomes in people with diabetes.

"Poor sleep quality in people with diabetes was associated with worse control of their blood glucose levels," said Kristen Knutson, PhD, assistant professor of medicine and lead author of the study. "People who have a hard time controlling their blood glucose levels have a greater risk of complications. They have a reduced quality of life. And, they have a reduced [life expectancy](#)."

People with diabetes generally have poorer sleep than the general population, and poor sleep has been proposed as a risk factor for developing the disease. [Sleep disorders](#), such as obstructive sleep apnea, are more prevalent in people with [type 2 diabetes](#), Knutson said.

For the study, researchers monitored the sleep of 40 people with diabetes for six nights. The subjects also reported if they generally suffered from symptoms of [sleep disturbances](#) like [insomnia](#), snoring or sleep apnea. At clinical examinations, they gave blood samples to allow researchers to measure insulin and glucose levels.

The subjects wore activity monitors on their wrists at night, which

measure their wrist movements throughout the night. Poor sleep, or insomnia, was determined by both poor sleep quality based on the activity monitors and the subject telling the researchers that they often had a hard time falling asleep or woke up during the night.

Among the diabetics, poor sleepers had 23% higher [blood glucose levels](#) in the morning, and 48% higher blood insulin levels. Using these numbers to estimate a person's [insulin resistance](#), the researchers found that poor sleepers with diabetes had 82% higher insulin resistance than normal sleepers with diabetes.

Knutson said the next step for researchers is to see if treating poor sleep can improve long-term outcomes and quality of life for diabetics. "For someone who already has diabetes, adding a sleep treatment intervention, whether it's treating sleep apnea or treating insomnia, may be an additional help for them to control their disease," Knutson said.

In fact, restoring a healthy amount of sleep may be as powerful an intervention as the drugs currently used to treat type 2 diabetes. "This suggests that improving sleep quality in diabetics would have a similar beneficial effect as the most commonly used anti-diabetes drugs," said Eve Van Cauter, PhD, professor of medicine and co-author of the study.

Further investigation into which leads to the other – the chronic poor sleep or chronic insulin resistance – could improve the quality of life for people with type 2 diabetes. "Anything that we can do to help people improve their ability to control their glucose will help their lives in the long run," Knutson said.

The data was collected as part of the CARDIA study, an ongoing longitudinal study of the heart health. It has tracked thousands of people for over 20 years.

**More information:** The study, "Cross-sectional associations between measure of sleep and markers of glucose metabolism among persons with and without diabetes" was published online March 16, 2011, in the journal *Diabetes Care* and is freely available.

Provided by University of Chicago Medical Center

Citation: Insomnia linked to high insulin resistance in diabetics (2011, May 2) retrieved 5 May 2024 from <https://medicalxpress.com/news/2011-05-insomnia-linked-high-insulin-resistance.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--