

27 percent of children wait too long for surgery

May 2 2011

Twenty-seven percent of children in Canada awaiting surgery at pediatric hospitals received the procedures past the target date, found an article in *CMAJ* (*Canadian Medical Association Journal*).

Prolonged wait times are a pressing issue in health care systems and Canadian governments at the federal, provincial and territorial levels have made reducing wait times a priority. However, some procedures may need to be performed sooner than by the established six-month target date.

The study, by researchers from the Hospital for Sick Children (SickKids) and the Canadian Paediatric Surgical Wait Times Project, was based on 64 012 surgeries at Canada's pediatric academic health centres. Dentistry (45%), ophthalmology (43%) and plastic surgery (35%) had the highest percentage of delayed surgeries followed by cancer, neurosurgery, and [cardiac surgery](#).

"Despite receipt of surgery within six months of the decision to treat for 90% of patients, clinically derived targets found that 27% of children waited too long for their specific condition," writes Dr. James G. Wright, The Hospital for Sick Children, Department of Surgery, with coauthors.

Delays in surgery can affect children's physical and [cognitive development](#). For example, delaying surgery to correct strabismus risks a child's chances of having normal vision and can affect quality of life.

"We believe this national collaborative project shows the feasibility and potential benefits of setting national standards for surgical wait times. Such accountability can result in measurably improved access in the delivery of national publicly funded health care," they conclude.

In a related research article, researchers found that waiting for surgery beyond wait time targets can affect the outcomes and risks associated with surgery. The authors found that wait times longer than three months for adolescents with scoliosis increase the likelihood of complications and additional surgeries. Surgery to correct idiopathic scoliosis, an abnormal curvature of the spine that is potentially progressive during adolescence, is the most common reason for elective pediatric orthopedic surgery.

"The highest risks of adverse events due to prolonged wait times occurred in patients who were skeletally immature and had larger curvatures of the spine," writes author Dr. Henry Ahn, St. Michael's Hospital, Toronto, with coauthors.

They conclude that a three-month wait time, based on empirical analysis rather than on conventional methods such as expert opinion, helps minimize risk. "A maximal acceptable wait time that leads to a reduction in the progression of curvature also has the potential to reduce the strain on healthcare resources by decreasing the need for additional [surgery](#)."

More information: www.cmaj.ca/cgi/doi/10.1503/cmaj.101530

Provided by Canadian Medical Association Journal

Citation: 27 percent of children wait too long for surgery (2011, May 2) retrieved 27 April 2024 from <https://medicalxpress.com/news/2011-05-percent-children-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.