

Ultrasound combined with pre-operative MRI is cost effective in evaluating rotator cuff tears

May 5 2011

While ultrasound is usually viewed as more cost effective, MRI is most often used to evaluate the rotator cuff. When performing a cost utility analysis, utilizing an ultrasound as the initial imaging test for rotator cuff tear, along with pre-operative MRI to identify alternative and concurrent diagnoses, can be a very effective hybrid imaging strategy, according to research being presented at the 2011 American Roentgen Ray Society's annual meeting.

The study, performed at Duke University Medical Center in Durham, NC and Rush University Medical Center in Chicago, IL, utilized three evaluation techniques: the use of [ultrasound](#) alone, MRI alone, and a hybrid strategy of ultrasound for all patients followed by MRI for those patients who required surgery.

"Several meta-analyses in the literature have found that ultrasound and MRI have similar accuracies for the evaluation of rotator cuff tears. Ultrasound is a cheaper imaging modality, yet MRI is much more frequently used for rotator cuff evaluation," said Robert Lee Suber, MD, lead author of the study. "The reasons for the preference of MRI may be related to the possibility of identifying alternative and/or concurrent diagnoses with MRI as well as surgeon preference for anatomic imaging prior to surgery," said Suber.

"One of the imaging strategies we studied was an initial screening test

with ultrasound. All those patients who required surgery or failed conservative treatment would then have an MRI. We found this to more cost effective than everyone undergoing MRI as the initial evaluation," he said.

"There are cutoff values for accuracy of ultrasound and/or MRI where this combined imaging strategy is no longer cost effective over MRI alone. Additionally, as the prevalence (pre-test probability) for rotator cuff tear increases, this combined imaging strategy decreases in cost savings over MRI alone," said Suber.

"Our research shows that in populations with a lower pre-test probability of rotator cuff tear (e.g. patients seeing family practice physicians as opposed to a shoulder specialist surgeon) it may be more cost effective to initially to obtain an ultrasound. Then if the patient needs to have surgery, they can get an [MRI](#)," he said.

This abstract is being presented in conjunction with the 2011 American Roentgen Ray Society's annual meeting.

Provided by American Roentgen Ray Society

Citation: Ultrasound combined with pre-operative MRI is cost effective in evaluating rotator cuff tears (2011, May 5) retrieved 2 May 2024 from <https://medicalxpress.com/news/2011-05-ultrasound-combined-pre-operative-mri-effective.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--