

# Brain structure adapts to environmental change

June 13 2011

---

Scientists have known for years that neurogenesis takes place throughout adulthood in the hippocampus of the mammalian brain. Now Columbia researchers have found that under stressful conditions, neural stem cells in the adult hippocampus can produce not only neurons, but also new stem cells. The brain stockpiles the neural stem cells, which later may produce neurons when conditions become favorable. This response to environmental conditions represents a novel form of brain plasticity. The findings were published online in *Neuron* on June 9, 2011.

The hippocampus is involved in memory, learning, and emotion. A research team led by Alex Dranovsky, MD, PhD, assistant professor of clinical psychiatry at Columbia University Medical Center and research scientist in the Division of Integrative Neuroscience at the New York State Psychiatric Institute/Columbia Psychiatry, compared the generation of neural stem cells and [neurons](#) in mice housed in isolation and in mice housed in enriched environments. They then used lineage studies, a technique that traces stem cells from their formation to their eventual differentiation into specific cell types, to see what proportion of neural stem cells produced neurons.

Deprived and enriched environments had opposite effects. The brains of the socially isolated mice accumulated neural stem cells but not neurons. The brains of mice housed in enriched environments produced far more neurons, but not more stem cells. The average mouse [dentate gyrus](#), the area of the hippocampus where neurogenesis takes place, has about 500,000 neurons; the enriched environment caused an increase of about

70,000 neurons.

"We already knew that enriching environments are neurogenic, but ours is the first report that neural stem cells, currently thought of as 'quiescent,' can accumulate in the live animal," said Dr. Dranovsky.

"Since this was revealed simply by changing the animal's living conditions, we think that it is an adaptation to stressful environments. When conditions turn more favorable, the stockpiled stem cells have the opportunity to produce more neurons—a form of 'neurons on demand.'"

The researchers also looked at neuronal survival. They found that social isolation did not cause it to decrease. Scientists already knew that environmental enrichment increased neuronal survival—further increasing the neuron population.

To a lesser extent, location within the [hippocampus](#) affected whether stem cells became neurons. While the ratio of stem cells to neurons remained constant in the lower blade of the dentrate gyrus, it varied in the upper blade.

Age also affected the results. After three months, the brains of the isolated mice stopped accumulating neural stem cells. But the mice in enriched environments continued to produce more neurons.

Dranovsky and his team now want to see whether this hippocampal response is specific to social isolation or is a more general response to stress. Another question is whether all neural stem cells have the same potential to produce neurons.

"The long-term goal." Said Dr. Dranovsky, "is to figure out how to instruct [neural stem cells](#) to produce neurons or more stem cells. This could lead to the eventual use of stem cells in neuronal replacement therapy for neurodegenerative diseases and other central nervous system

conditions."

Provided by Columbia University

Citation: Brain structure adapts to environmental change (2011, June 13) retrieved 30 April 2024 from <https://medicalxpress.com/news/2011-06-brain-environmental.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.