

Childhood cancer survivors at increased risk of certain tumors in middle-age

June 5 2011

In a study that included nearly 18,000 children who had cancer, with follow-up of about 25 years, the greatest excess risk associated with a subsequent primary neoplasm (a new tumor) at older than age 40 years was for digestive and genitourinary neoplasms (related to the genital or urinary tract organs), according to a study in the June 8 issue of *JAMA*, a theme issue on cancer.

"Survivors of <u>childhood cancer</u> are at increased risk of developing subsequent primary <u>neoplasms</u> with the risk being estimated at 3- to 6-fold than expected. However, the magnitude of the long-term risks of developing a subsequent primary neoplasm, in particular beyond ages at which the background <u>cancer incidence</u> in the general population starts to increase substantially, are unknown," according to background information in the article. "It is essential to investigate the long-term risks of subsequent primary neoplasms since any increased relative risk (RR) sustained into old age would lead to a substantial number of survivors being diagnosed with a subsequent primary neoplasm. To reduce the number of subsequent primary neoplasms, prevention, screening, and other intervention strategies should focus on subsequent primary neoplasm types with the highest absolute excess risks (AERs); however, it is currently unclear which subsequent primary neoplasm types contribute most to the total AER, particularly at older than 40 years."

Raoul C. Reulen, Ph.D., of the University of Birmingham, Edgbaston, United Kingdom, and colleagues conducted a study to investigate the



long-term risks of subsequent primary neoplasms in survivors of childhood cancer. The British Childhood Cancer Survivor Study included 17,981 5-year survivors of childhood cancer diagnosed with cancer at younger than 15 years between 1940 and 1991 in Great Britain, who were followed up through December 2006.

Through the follow-up period, 1,354 subsequent primary neoplasms were diagnosed in 1,222 of the 17,981 survivors in the cohort. The average follow-up time from childhood cancer diagnosis was 25.6 years. The most commonly observed subsequent primary neoplasms were central nervous system tumors (n = 344) of which 105 were gliomas, nonmelanoma skin cancer (n = 278), digestive (n = 105), genitourinary (n = 100), breast (n = 97), and bone (n = 94). Overall, survivors were 4 times more likely to develop a subsequent primary neoplasm than expected. Digestive and genitourinary subsequent primary neoplasms contributed the largest AER at older than 40 years compared with other subsequent primary neoplasm categories; 36 percent of the total AER was attributable to genitourinary and digestive subsequent primary neoplasms. Overall, 52 percent of the total AER was attributable to digestive, genitourinary, breast, or respiratory sites at older than 40 years.

"These increased risks of digestive subsequent primary neoplasms are likely to be related to previous exposure of the digestive tract to radiation, except following heritable retinoblastoma. We found a 3-fold increased RR for survivors treated with direct abdomino-pelvic irradiation. The highest AER for genitourinary subsequent primary neoplasms was observed after heritable retinoblastoma, suggesting that this excess risk is at least partially attributable to genetic predisposition," the authors write.

The cumulative percentage of survivors developing a subsequent primary neoplasm increased steadily with attained age from 1.6 percent at age 20



years to 13.8 percent at age 60 years, whereas 8.4 percent was expected at the latter age based on rates from the general population. Five percent of survivors had developed a subsequent primary neoplasm by age 38 years, whereas it took until age 54 years for 5 percent of a comparable cohort (based on the general population rates) to develop a cancer.

The researchers also found that the cumulative incidence of developing colorectal cancer by age 50 years was 1.4 percent for survivors treated with direct abdomino-pelvic irradiation. This risk is comparable to the equivalent percentage (1.2 percent) for individuals with at least 2 first-degree relatives affected by colorectal cancer. They add that this is, to their knowledge, the first time this finding has been demonstrated. "This raises the question of whether childhood <u>cancer survivors</u> treated with direct abdomino-pelvic irradiation should be similarly considered for routine colonoscopy for colorectal cancer."

The authors note that it should be emphasized that the AER for most subsequent primary neoplasms was still relatively low; although for genitourinary and digestive subsequent primary neoplasms, it increased rapidly with attained age, and if continued into old age, a substantial number of survivors would develop such subsequent primary neoplasms. "Given the excesses observed among those older than 40 years, these survivors should be encouraged to participate in existing general population screening programs."

More information: JAMA. 2011;305[22]2311-2319

Provided by JAMA and Archives Journals

Citation: Childhood cancer survivors at increased risk of certain tumors in middle-age (2011, June 5) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2011-06-childhood-cancer-</u>



survivors-tumors-middle-age.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.