

# Irregular breathing can affect accuracy of 4-D PET/CT

June 7 2011

---

A study presented at SNM's 58th Annual Meeting focuses on the effect that breathing irregularities have on the accuracy of 4D positron emission tomography (PET) scans and outlines a PET imaging method that reduces "motion artifacts" or image blurring arising from respiratory motion. Non-gated PET imaging with 4D computed tomography may be useful for imaging patients who do not benefit from the use of respiratory gating, most notably patients with erratic breathing.

"Breathing irregularities can lead to significantly underestimated lesion activity in respiratory-gated [PET imaging](#)," says Boon-Keng Teo, PhD, assistant professor of [radiation oncology](#) at the University of Pennsylvania in Philadelphia, Pa. "Non-gated PET imaging corrected with 4D computed tomography (CT) may be more effective for imaging patients with irregular breathing. This could potentially lead to a more robust and quantitatively accurate reading of active tumors."

Respiratory gating technologies have dramatically improved the diagnostic quality of PET imaging, which provides functional images of [physiological processes](#) occurring in the body. Sensors in respiratory gating systems placed on or around the patient monitor the phase of the breathing cycle. They then transmit information about the patient's breathing to the [scanning technology](#) for image processing. Instead of creating one fluid image that shows so-called motion artifacts, respiratory-gated PET imaging is much like a series of photos taken during different phases of the respiratory cycle that are grouped together to create a series of images corresponding to each phase. The problem is

that patients with respiratory disease, [heart conditions](#) or other serious disease are likely to be breathing unevenly. Respiratory gating systems are designed to work with normal breathing patterns, but not with irregular respiratory cycles.

Researchers conducted phantom studies to compare respiratory-gated PET imaging with non-gated PET imaging corrected with 4D computed tomography. CT uses X-ray technology and complex data processing to produce very high-resolution images of structural anatomy. Phantom studies were performed with inanimate objects and specialized motion devices that move in a controlled manner in order to simulate tumors in respiratory motion. The 4D PET and CT studies were conducted in succession with a hybrid PET/CT system. Various degrees of motion irregularities were simulated to study their impact on the accuracy of 4D PET for suppressing motion artifacts.

Results of the study show that non-gated PET with 4D CT imaging can be an alternative to respiratory-gated PET imaging for determining tumor activity in patients with highly irregular breathing. These findings could change imaging protocols for patients with uneven breathing and potentially improve overall accuracy of tumor detection, thereby informing clinicians about appropriate treatments and perhaps even surgical planning for better cancer management.

**More information:** Scientific Paper 149: B. Teo, B. Saboury, J. Scheuermann, D. Torigian, J. Karp, A. Alavi, University of Pennsylvania, Philadelphia, PA; "Effect of breathing irregularities on the quantitative accuracy of respiratory gated PET/CT," SNM's 58th Annual Meeting, June 4-8, 2011, San Antonio, TX.

Provided by Society of Nuclear Medicine

Citation: Irregular breathing can affect accuracy of 4-D PET/CT (2011, June 7) retrieved 5 May 2024 from <https://medicalxpress.com/news/2011-06-irregular-affect-accuracy-d-petct.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.