

Molecular imaging detects ischemic heart disease in diabetics

June 6 2011

Research introduced at SNM's 58th Annual Meeting may lead to much-needed cardiovascular disease screening for diabetic patients at risk of ischemic heart disease, a disorder marked by significantly reduced blood flow in the heart. Ischemia of the myocardium, or cardiac muscle, can signal diminished oxygenation of the heart tissue and trigger a heart attack if left untreated.

"Diabetes is a serious risk factor for ischemic heart disease," says Kenji Fukushima, MD, PhD, Tokyo Women's Medical University, Tokyo, Japan. "Most [diabetic patients](#) do not exhibit signs of [myocardial ischemia](#) and may be on course toward major cardiac events without any red alert. [Molecular imaging](#) could potentially provide an effective [screening tool](#) for at-risk diabetic patients suspected of having ischemic heart disease."

According to the American Diabetes Association, approximately 25.8 million children and adults—more than eight percent of the population—are diabetic and at least twice as likely as non-diabetics to develop heart disease. The link is attributed to the damaging effects of high blood sugar levels on blood vessels. Persistently high blood sugar levels can lead to inflammatory disease and atherosclerosis, or thickening of the blood vessel walls, and eventually myocardial ischemia.

The standard molecular imaging technique is stress myocardial perfusion imaging, which provides information about blood flow in the heart at rest and under stress either with exercise or with drugs that recreate the

physiological changes associated with exercise. However, stress testing may be unsafe for patients at risk of complications or cardiac events. This study explores a method of molecular imaging called myocardial fatty acid metabolism imaging, which may prove to be a safer alternative for patients suspected of having ischemic heart disease.

During the study, a total of 191 diabetic patients were injected with I-123 beta-methyl iodophenyl pentadecanoic acid (BMIPP), a medical isotope bound with an agent that is metabolized by the heart in order to image the fatty acid uptake of heart muscle cells. Once injected, patients were imaged using dual single photon emission computed tomography (SPECT), a molecular imaging technology that captures both physiological and anatomical information about the body. The objective of the study was to test the predictive value of this imaging procedure for major cardiac events like fatal heart attack and minor cardiac events such as hospitalization due to heart failure. Follow-up of the study revealed 13 major and 25 minor events, proving that I-123 BMIPP is safe and beneficial for predicting the likelihood of [cardiac events](#) in diabetic patients suspected of having myocardial ischemia.

This is the first study of its kind detecting ischemic heart disease in diabetic patients without any history of myocardial disease. If moved into clinical use, imaging both cell metabolism and blood flow through the heart could help physicians detect myocardial ischemia and make clinical decisions that could save lives.

More information: Scientific Paper 110: K. Fukushima, M. Momose, Y. Miyake, C. Kondo, T. Nakajima, N. Hagiwara, S. Sakai, Tokyo Women's Medical University, Tokyo, Japan; "Prognostic usefulness of I-123 BMIPP and TL-201 dual SPECT in patients with diabetes mellitus for suspected ischemic heart disease," SNM's 58th Annual Meeting, June 4-8, 2011, San Antonio, TX.

Provided by Society of Nuclear Medicine

Citation: Molecular imaging detects ischemic heart disease in diabetics (2011, June 6) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2011-06-molecular-imaging-ischemic-heart-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.