

## Sleepiness may impair the brain's inhibitory control when viewing high-calorie foods

June 13 2011

Daytime sleepiness may affect inhibitory control in the brain when viewing tantalizing, high-calorie foods, suggests a research abstract that will be presented Monday, June 13, in Minneapolis, Minn., at SLEEP 2011, the 25th Anniversary Meeting of the Associated Professional Sleep Societies LLC (APSS).

Results show that greater <u>daytime sleepiness</u> was associated with decreased activation in the prefrontal cortex during visual presentations of enticing, high-calorie food images. The prefrontal cortex is a <u>brain</u> region that plays an important role in inhibitory processing.

"Self-reported daytime sleepiness among healthy, normally rested individuals correlated with reduced responsiveness of inhibitory brain regions when confronted with images of highly appetizing foods," said principal investigator William Killgore, PhD, assistant professor of psychology at Harvard Medical School and McLean Hospital in Belmont, Mass. "It suggests that even normal fluctuations in sleepiness may be capable of altering brain responses that are important for regulating dietary intake, potentially affecting the types of choices that individuals make when selecting whether and what to eat."

The research team of Killgore, lead author Melissa Weiner, and Zachary Schwab studied 12 healthy men and women between the ages of 19 and 45 years. The participants underwent functional <u>magnetic resonance</u> imaging (fMRI) while viewing pictures of high-calorie foods, low-calorie foods, and control images of plants and rocks. Subjective, self-



reported daytime sleepiness was measured with the Epworth Sleepiness Scale, which evaluates how likely an individual is to doze off or fall asleep during certain situations such as while sitting and reading or watching TV.

According to the authors, prior evidence suggests that healthy adults activate inhibitory regions of the <u>prefrontal cortex</u> in response to high-calorie food images. However, insufficient sleep is often associated with reduced metabolic activity within these same prefrontal regions.

Killgore noted that the rapidly rising rate of obesity makes it important to understand the relationship between sleep-related factors, brain responses to food, and eating behavior.

"Given the chronic level of sleep restriction in our society, such relationships could have epidemiologic implications regarding the current increase in obesity in westernized countries," he said.

In a previous study published in Neuroreport in 2010, Killgore also found sex differences in cerebral responses to the caloric content of food images. Results of that study indicate that when viewing high-calorie food images, women showed significantly greater activation than men in brain regions that are involved in behavioral control and self-referential cognition.

## Provided by American Academy of Sleep Medicine

Citation: Sleepiness may impair the brain's inhibitory control when viewing high-calorie foods (2011, June 13) retrieved 27 June 2024 from <u>https://medicalxpress.com/news/2011-06-sleepiness-impair-brain-inhibitory-viewing.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.