

# Surgery deaths drop nationwide for high-risk surgeries: study

June 1 2011

---



In the past decade, surgery death rates have dropped dramatically for even the most complicated surgeries. But a University of Michigan Health System study in this week's *New England Journal of Medicine* reveals the decline is linked with more surgeries shifting to safer, high-volume hospitals, particularly surgeries to treat pancreatic and esophageal cancers. Credit: University of Michigan Health System

Surgery death rates have dropped nationwide over the past decade, according to a University of Michigan Health System study that reveals cancer surgeries have seen the most dramatic improvement in safety.

The U-M study in this week's [New England Journal of Medicine](#) shows surgery mortality dropped substantially for eight different high-risk surgeries performed on 3.2 million [Medicare patients](#) from 1999 to 2008.

More patients are surviving [open heart surgery](#) and replacement of diseased aortic valves, but research shows high volume hospitals and their expertise drove a 67 percent decline in deaths for pancreatectomy, a 37 percent decline in deaths from cystectomy, surgery to remove the bladder, and a 32 percent drop in esophagectomy mortality.

"Patients should take solace in knowing that all high-risk surgeries have become safer in the last decade," says lead author Jonathan F. Finks, M.D., clinical assistant professor of surgery at the U-M Health System. "In [cancer surgery](#), in particular, mortality has dropped in large part because more patients are having their surgery in safer, higher volume hospitals."

One of the more interesting findings by the U-M's Center for Healthcare Outcomes and Policy is that hundreds of low-volume U.S. hospitals stopped doing high-risk cancer surgery.

For example, the number of Medicare patients needing surgery to treat pancreatic cancer increased by 50 percent, but the number of hospitals performing the surgeries decreased by 25 percent, from 1,308 hospitals to 978.

As a result, the volume of surgeries a hospital may do in a year rose from five cases of pancreatectomies a year to 16.

There have been numerous efforts in the United States to concentrate selected operations at high volume hospitals.

The Leapfrog group, a consortium of large corporations and public agendas that purchase health care largely for their employees, has been among the most prominent advocates of volume-based referrals. In 2000, it placed a minimum volume standard on hospitals for several surgical procedures to guide where employees get medical care.

The U-M analysis shows volume-based referrals have been successful for certain surgeries, but the effort to make surgery safer takes more than one fix, authors say.

"For some procedures, however, strategies such as operating room checklists, outcomes measurement programs, and quality improvement collaboratives are likely to be more effective than volume-based referral," says John Birkmeyer, M.D., professor of [surgery](#) and director of the U-M Center for Healthcare Outcomes and Policy.

Provided by University of Michigan

Citation: Surgery deaths drop nationwide for high-risk surgeries: study (2011, June 1) retrieved 4 May 2024 from <https://medicalxpress.com/news/2011-06-surgery-deaths-nationwide-high-risk-surgeries.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--