

Harmful effects of hypothyroidism on maternal and fetal health drive new guidelines for managing thyroid disease in preg

July 25 2011

Emerging data clarifying the risks of insufficient thyroid activity during pregnancy on the health of the mother and fetus, and on the future intellectual development of the child, have led to new clinical guidelines for diagnosing and managing thyroid disease during this critical period. The guidelines, developed by an American Thyroid Association (ATA) expert task force, are presented in *Thyroid*.

Clinical studies are producing critical data demonstrating the harmful effects not only of overt [hypothyroidism](#) and [hyperthyroidism](#) on pregnancy, but also of subclinical [thyroid](#) disease and maternal and fetal health. Ongoing research is clarifying the link between miscarriage and preterm delivery in women with normal [thyroid function](#) who are thyroid peroxidase antibody positive. Studies are also uncovering the long-term effects of postpartum thyroiditis.

"Pregnancy has a profound impact on the [thyroid gland](#) and thyroid function.... In essence, pregnancy is a stress test for the thyroid, resulting in hypothyroidism in women with limited thyroidal reserve or iodine deficiency," state Alex Stagnaro-Green, George Washington University School of Medicine and Health Sciences (Washington, DC), and coauthors representing the ATA task force.

Among the many specific recommendations detailed in the guidelines

are the following: women with overt hypothyroidism or with subclinical hypothyroidism who are TPO antibody positive should be treated with oral levothyroxine; use of other thyroid preparations such as triiodothyronine or desiccated thyroid to treat maternal hypothyroidism is strongly recommended against; and women with subclinical hypothyroidism in pregnancy who are not initially treated should be monitored for progression to overt hypothyroidism with serum TSH and free T4 measurements about every 4 weeks until 16-20 weeks gestation and at least once between 26-32 weeks gestation.

The new clinical guidelines focus on several key areas in the diagnosis and management of thyroid disease during pregnancy and postpartum: thyroid function tests, hypothyroidism, thyrotoxicosis, iodine, thyroid antibodies and miscarriage/[preterm delivery](#), thyroid nodules and cancer, postpartum thyroiditis, recommendations on screening for thyroid disease during pregnancy, and areas for future research.

"These important guidelines were developed by a panel of international experts representing the disciplines of endocrinology, obstetrics and gynecology, and nurse midwives. This broad representation of providers that care for pregnant women will significantly increase the impact of these guidelines and translation of findings from the most recent research to clinical practice," says Gregory A. Brent, MD, Professor of Medicine and Physiology, David Geffen School of Medicine at the University of California Los Angeles and President of the ATA.

"[Thyroid disease](#) in pregnancy is common, clinically important, and time-sensitive, and our knowledge about it is rapidly changing. This ATA Guideline will disseminate this new information both widely and rapidly to improve patient care, establish what we believe is optimal care for the pregnant woman and her unborn child, and drive future research to further improve our understanding and patient outcomes," says Richard T. Kloos, MD, Professor, The Ohio State University and Secretary/Chief

Operating Officer of the ATA.

More information: www.liebertpub.com/thy

Provided by Mary Ann Liebert, Inc.

Citation: Harmful effects of hypothyroidism on maternal and fetal health drive new guidelines for managing thyroid disease in preg (2011, July 25) retrieved 4 May 2024 from <https://medicalxpress.com/news/2011-07-effects-hypothyroidism-maternal-fetal-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.