

Study examines factors in pediatric kidney transplant rejection

July 18 2011

Avoiding HLA-DR mismatching appears to be beneficial in pediatric kidney transplant patients, however the likelihood of finding a matching donor must be considered against the wait time for a possible donation, according to a report in the July issue of *Archives of Surgery*.

"Although avoiding HLA [human leukocyte antigen; cell surface antigens that regulate [host cell](#) responses to transplanted cells] antigen mismatching has been shown to benefit long-term graft survival, it has raised concerns about disadvantaging minority groups, particularly black patients, and [pediatric patients](#), who have severe growth retardation and other problems when dialysis is prolonged before transplantation," the authors write as background information in the article. "Currently, only HLA-DR matching is considered in the United Network for Organ Sharing (UNOS) organ allocation system."

To examine the relationship between HLA-DR mismatching and rejection, graft survival and sensitization in pediatric [kidney transplant patients](#), Lan T. Vu, M.D., and colleagues from the University of California at San Francisco, conducted a [retrospective cohort study](#) of 178 pediatric patients who underwent primary [kidney transplantation](#) with daclizumab induction therapy (to prevent [organ rejection](#)) at the University of California, San Francisco between 1997 and 2006.

One year after transplantation, 35 percent of the patients experienced rejection and at five year follow-up the frequency of rejection was 55 percent. Patients with 1- or 2-HLA-DRB1 mismatches had 1.7 times

greater odds of rejection than patients with no HLA-DRB1 mismatches. "This single-center study demonstrated that HLA-DRB1 mismatching increased the risk of allograft rejection by approximately 70 percent in children," write the authors.

Additionally, the 1- and 5-year graft survival rates for this study group were 97 percent and 82 percent respectively, and the authors found that the degree of HLA-DRB1 mismatches were not significantly related to graft failure. However, patients with a history of rejection had 7.7 times greater odds of graft failure than those who had not previously had an episode of rejection.

"In conclusion, this study showed that HLA-DRB1 mismatch was a risk factor for rejection and that rejection was a strong predictor of graft failure and sensitization in children," the authors write. "Based on the high incidence of rejection and sensitization seen with HLA-DR-mismatched kidneys observed in this series, we advocate expansion of the current artificial boundaries of donor pools to facilitate better matching and outcomes for young recipients."

More information: Arch Surg. 2011;146[7]:824-829.

Provided by JAMA and Archives Journals

Citation: Study examines factors in pediatric kidney transplant rejection (2011, July 18) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2011-07-factors-pediatric-kidney-transplant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.