

Genetic study sheds new light on autoimmune arthritis

July 10 2011

The team of researchers from the Universities of Bristol, Queensland. Oxford, Texas and Toronto, used a technique called genome-wide association where millions of genetic markers are measured in thousands of people that have the disease and thousands of healthy individuals.

Markers which are more frequent in individuals with the disease are more likely to be involved in the condition.

Using this approach the investigators found an additional seven genes likely to be involved in the condition, bringing the total number of genes known to predispose to AS to thirteen. Many of the new genes are already known to be involved in inflammatory and immune processes, providing researchers with further clues about how the disease arises. Two of the new genes are also known to predispose to other autoimmune conditions including <u>Crohn's disease</u> (a form of <u>inflammatory</u> <u>bowel disease</u>) and <u>Celiac disease</u> (an auto-immune intestinal disease).

Researchers were also able to demonstrate an interaction between a genetic mutation called HLA-B27 and a mutation in a gene called ERAP1. Specifically, the ERAP1 mutation only predisposed to disease in those individuals who tested positive for the HLA-B27 mutation.

Dr David Evans from the University of Bristol said: "This finding is important in a number of ways. First of all it's one of the first convincing examples we have of one mutation influencing the effect of another mutation in the development of a relatively common disease. This is



exciting because it implies that there may be other examples of this phenomenon in other <u>common diseases</u> that we don't know about yet.

"Second, the interaction itself tells us something very fundamental about how AS is caused. Prior to this study there were a number of competing theories about how the disease was caused. Our study suggests very strongly which one of these competing <u>hypotheses</u> is likely to be correct."

Finally, the researchers also identified a single genetic marker which could be used to assist in diagnosis of Ankylosing Spondylitis.

"AS is notoriously difficult to diagnose in its early stages which can lead to costly delays in its treatment," said Dr Evans. "Typically diagnosis consists of a combination of X-rays, patient symptoms and expensive immunological assays in the laboratory. This <u>genetic marker</u> could easily take the place of an expensive immunological assay. What would normally cost £40-£50 could be done easily for a fraction of the price."

More information: 'Interaction between ERAP1 and HLA-B27 in ankylosing spondylitis implicates peptide handling as the mechanism for HLA-B27 in disease susceptibility', Evans et al. *Nature Genetics*, July 10, 2011.

Provided by University of Bristol

Citation: Genetic study sheds new light on auto-immune arthritis (2011, July 10) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2011-07-genetic-auto-immune-arthritis.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.