

Hearing loss is another danger of secondhand smoke

July 18 2011

NYU School of Medicine researchers report in a new study that exposure to tobacco smoke nearly doubles the risk of hearing loss among adolescents. The study is published in the July, 2011, issue of *Archives of Otolaryngology – Head & Neck Surgery*.

"More than half of all children in the U.S. are exposed to secondhand smoke, so our finding that it can lead to [hearing loss](#) in teenagers has huge public health implications,"* says Anil Lalwani, MD, professor of professor of otolaryngology, physiology and neuroscience, and pediatrics at NYU School of Medicine, who led the research. "We need to evaluate how we deal with smoking in public places and at home, as well as how often and when we screen children for hearing loss," he says.

The dangers of secondhand smoke are well known. Living with a smoker raises the risk of dying from heart disease and lung cancer, and in children exposure to smoke exacerbates the severity of asthma attacks and causes more than 750,000 middle ear infections, according to the American Cancer Society. The new study is the first to link secondhand smoke to hearing loss.

More than 1,500 teenagers aged 12 to 19 participated in the nationwide study. They were selected from the 2005-2006 National Health and Nutrition Examination Survey, which collects health information from children and adults around the United States. The teenagers were initially evaluated in their homes and then were given extensive hearing tests and blood tests for the chemical cotinine, a metabolite of nicotine, at a

medical center.

The teens exposed to secondhand smoke, as measured by the metabolite in their blood, were more likely to have sensorineural hearing loss, which is most often caused by problems with the cochlea, the snail-shaped hearing organ of the inner ear. "It's the type of hearing loss that usually tends to occur as one gets older, or among children born with congenital deafness," explains co-author Michael Weitzman, MD, professor of pediatrics and psychiatry at NYU School of Medicine.

The study found that teenagers exposed to smoke performed worse across every sound frequency tested, especially mid-to-high frequencies important for understanding speech. In addition, teenagers with higher cotinine levels, indicating greater exposure, were more likely to have one-sided—or unilateral—low-frequency hearing loss. Overall, the researchers conclude that their findings indicate that "[tobacco smoke](#) is independently associated with an almost 2-fold increase in the risk of hearing loss among [adolescents](#)."

Over 80 percent of the affected teenagers in the study were not aware of any problem, the researchers reported. "Milder hearing loss is not necessarily noticeable," says Dr. Lalwani. "Thus, simply asking someone whether they think they have hearing loss is insufficient."

The consequences of mild hearing loss, which researchers suspect may be due to damage to the ear's delicate blood supply, are "subtle yet serious," says Dr. Weitzman. Affected children can have difficulty understanding what is being said in the classroom and become distracted. As a result, they may be labeled as "troublemakers" or misdiagnosed with ADHD (attention deficit hyperactivity disorder).

Currently, all infants born in the United States are screened for hearing loss; however, there are no guidelines for screening a child's hearing past

the early school years, says Dr. Lalwani. "Those [children](#) who are exposed to [secondhand smoke](#)," he says, "need to be regularly screened."

Provided by New York University School of Medicine

Citation: Hearing loss is another danger of secondhand smoke (2011, July 18) retrieved 26 April 2024 from <https://medicalxpress.com/news/2011-07-loss-danger-secondhand.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.