

# 'Megapixel' DNA replication technology promises faster, more precise diagnostics

July 3 2011

---

UBC researchers have developed a DNA measurement platform that sets dramatic new performance standards in the sensitivity and accuracy of sample screening.

The advance could improve a range of genetic diagnostics and screenings where precise measurement is crucial--including the early detection of cancer, prenatal diagnostics, the detection of [pathogens](#) in [food products](#), and the analysis of single cell [gene expression](#).

The new digital [polymerase chain reaction](#) (PCR) device uses liquid [surface tension](#), rather than systems of microscopic valves, to partition [DNA samples](#) into arrays of 1,000,000 chambers or more. The device enables the direct counting of single molecules isolated in individual chambers.

The density of reaction chambers achieved by the platform exceeds more traditional valve-based digital PCR techniques by a factor of 100, translating directly into improved performance.

"This solves some major technical issues that have limited the scale and accuracy of traditional digital PCR techniques," says Assistant Professor Carl Hansen with the UBC Department of Physics and Astronomy and Centre for High-Throughput Biology. "It creates defect-free arrays of millions of uniform volume sub-reactions, and controls dehydration of these reactions during thermocycling."

PCR is an indispensable molecular biology technique used by researchers to amplify--or copy--a single piece of DNA millions or billions of times. The technique relies on repeated cycles of heating and cooling of the reaction to replicate segments of DNA using a protein called DNA polymerase, the same enzyme that copies DNA in living cells. PCR is used in medical and biology labs to clone DNA, analyze genes, detect hereditary disease, and in forensics.

The description of the 'megapixel' platform was published today in *Nature Methods*.

Digital PCR refers to a new generation of [DNA replication](#) techniques that offer increased sensitivity and density over the original technique, developed in 1983. The greatest number of chambers available in commercially available implementations of digital PCR, using integrated micro-valves, is 36,960. However, further scalability is limited by the maximum density at which valves may be reliably fabricated.

Hansen believes the new version of digital PCR can be scaled to hold up to approximately 10,000,000 chambers on a standard one inch format.

The UBC researchers also found the new 'megapixel' technique set new benchmarks in detecting rare mutations--defined as the lowest measurable ratio of two target sequences differing by a single nucleotide variation as well as new limits in the detection of subtle differences in sequence abundance.

Partitioning of a one million chamber array takes approximately one minute.

"Our solution, or something using the same techniques, could enable a new degree of precision in measurements in biomedical research and diagnostics. The dramatic increase in assay density has important

implications for the adoption of digital PCR as an economical, fast and routine analytical tool," says Hansen.

**More information:** Paper online: [dx.doi.org/10.1038/nmeth.1640](https://doi.org/10.1038/nmeth.1640)

Provided by University of British Columbia

Citation: 'Megapixel' DNA replication technology promises faster, more precise diagnostics (2011, July 3) retrieved 1 May 2024 from <https://medicalxpress.com/news/2011-07-megapixel-dna-replication-technology-faster.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--