

CABG still preferred over PCI in patients with triple vessel disease

August 29 2011

Results from CREDO-Kyoto PCI/CABG Registry Cohort-2 show that percutaneous coronary intervention (PCI) was associated with significantly higher risk for serious adverse events in patients with triple vessel disease than coronary artery bypass grafting (CABG). The protective effect of CABG for myocardial infarction was described as "especially remarkable". In view of these results, said investigator Dr Hiroki Shiomi from Kyoto University Hospital in Japan, "CABG would still remain the standard treatment option in patients with triple vessel disease, particularly when their SYNTAX scores are high".

These registry findings, derived from the largest ever study population of triple [vessel disease](#) patients with SYNTAX score assessment, are consistent with those found in the SYNTAX randomised trial. However, said Dr Shiomi, while the observations were striking in patients with triple vessel disease, the selection of revascularisation strategies in patients with less complex coronary anatomy "deserves further consideration".

As background to the report, he explained that PCI has been widely performed in patients with severe [coronary disease](#) (such as left main or triple vessel [coronary artery disease](#)) following the introduction of drug-eluting stents. However, long-term [clinical outcomes](#) of PCI relative to CABG in such patients have not yet been adequately evaluated. Although three-year results from the SYNTAX trial suggested that an excess risk of PCI relative to CABG for death, myocardial infarction or stroke was significant in the triple vessel disease subset, there were

limitations in the apparent lack of statistical power to evaluate this composite endpoint.

The CREDO-Kyoto (Coronary REvascularization Demonstrating Outcome Study in Kyoto) PCI/CABG registry cohort-2 is a physician-initiated non-company sponsored 26-centre registry enrolling consecutive patients having a first coronary revascularisation between January 2005 and December 2007. The study population for the current analysis consisted of 2981 patients with triple vessel disease (PCI 1825 patients, and CABG 1156 patients). To ensure comparability between the PCI and CABG groups (in an observational study), anatomic complexities of coronary artery disease were assessed by using the SYNTAX score.

The primary endpoint of the study was a composite of all-cause death, [myocardial infarction](#) (MI) and stroke. PCI as compared with CABG was associated with a higher 3-year risk for this primary endpoint (adjusted hazard ratio (HR) 1.47 [95% CI 1.13-1.92, P=0.004]) and for MI (HR 2.39 [95% CI 1.31-4.36, P=0.004]). However, the risk for cardiac death was not significantly different (HR 1.30 [95% CI 0.81-2.07, P=0.28]), although the risk for all-cause death was significantly higher after PCI (HR 1.62 [95% CI 1.16-2.27, P=0.005]).

Results also showed that the cumulative incidence of the primary endpoint was comparable between the PCI and CABG groups in patients with low (

Citation: CABG still preferred over PCI in patients with triple vessel disease (2011, August 29) retrieved 23 April 2024 from

<https://medicalxpress.com/news/2011-08-cabg-pci-patients-triple-vessel.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.