

Coronary artery stenting viable palliative option for infants and toddlers

August 22 2011

Percutaneous coronary intervention (PCI) is not a generally accepted option for infants or toddlers with acute coronary syndrome. However, a new report published in the August issue of *Catheterization and Cardiovascular Interventions*, a journal of The Society for Cardiovascular Angiography and Interventions (SCAI), found coronary stent implantation to be a feasible and safe palliative option for children fifteen months and younger.

Studies have shown that in adults with reduced blood flow to the heart—known as [acute coronary syndrome](#) (ACS)—[stent implantation](#) significantly reduces mortality. While ACS is less common in children there are congenital and acquired heart disease that may compromise coronary circulation in the pediatric population. Due to increased risk of coronary interventions and difficulty of cardiac catheterization techniques in pediatric patients, PCI is not a recommended treatment strategy in this age group.

A research team led by Dr. András Bratincsák from Rady Children's Hospital in San Diego, California, conducted a retrospective review of seven children who had PCI between June 2006 and June 2010. Of those included in the study all were under 18 years of age and four were 15 months or younger. PCI techniques included balloon coronary angioplasty and coronary stent implantation. Researchers analyzed patient data that included underlying diagnosis, comorbidities, catheterization technique, and outcomes.

In all seven cases, successful stent placement in the proximal portion of the left or right coronary arteries with excellent revascularization was achieved. The average diameter of the [heart](#) arteries was 0.65 mm prior to the intervention. Balloon angioplasty did not completely resolve the stenosis and bare metal stents were then implanted to a mean internal diameter of 2.5 mm. The team determined that the average intervention-free period was 434 days after stent implantation. Restenosis and thrombosis did not occur in cases where the implanted stent diameter was greater than 2.5 mm and patients received dual anti-platelet therapy.

"We provide evidence that stent implantation is a relatively safe option for pediatric patients with coronary stenosis, including those under the age of 15 months," said Dr. Bratincsák. "PCI offers a viable strategy for bridging infants and [toddlers](#) with blocked arteries or poor ventricular function to surgical revascularization or transplantation when they are at an older age."

More information: "Percutaneous Stent Implantation into Coronary Arteries in Infants." Andras Bratincsa, Anas Salkini, Howaida G. El-Said and John W. Moore. *Catheterization and Cardiovascular Interventions*; Published Online: August 22, 2011 ([DOI: 10.1002/ccd.23173](https://doi.org/10.1002/ccd.23173)); Print Issue Date: August 2011.

Provided by Wiley

Citation: Coronary artery stenting viable palliative option for infants and toddlers (2011, August 22) retrieved 24 April 2024 from <https://medicalxpress.com/news/2011-08-coronary-artery-stenting-viable-palliative.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.