

Diastolic dysfunction appears to worsen over time; associated with increased risk of heart failure

August 23 2011

A follow-up of participants in a heart function study finds that the prevalence of left ventricular diastolic dysfunction (left ventricular filling [with blood] is abnormal and is accompanied by elevated filling pressures) had increased; that diastolic function had worsened in a nearly a quarter of patients; and that participants who had diastolic dysfunction were more likely to develop heart failure, according to a study in the August 24/31 issue of *JAMA*.

"There is an emerging emphasis on understanding the progression from heart failure <u>risk factors</u> to asymptomatic ventricular dysfunction and eventually to symptomatic heart failure and death. Therefore, it is important to have population-based information on changes in <u>cardiac</u> <u>function</u> over time," according to background information in the article. "... little is known about time-dependent changes in diastolic function or their relationship to clinical heart failure."

Garvan C. Kane, M.D., Ph.D., of the Mayo Clinic and Medical School, Rochester, Minn., and colleagues conducted a study to measure changes in diastolic function over time and to determine the relationship between diastolic dysfunction and the risk of subsequent heart failure. The study included participants enrolled in the Olmsted County <u>Heart Function</u> Study (OCHFS). Randomly selected participants 45 years or older (n = 2,042) underwent clinical evaluation, medical record abstraction, and echocardiography (examination 1 [1997-2000]). Diastolic <u>left</u>



<u>ventricular</u> function was graded as normal, mild, moderate, or severe via testing. After 4 years, participants were invited to return for examination 2 (2001-2004). The group of participants returning for examination 2 (n = 1,402 of 1,960 surviving [72 percent]) then underwent follow-up testing for determination of new-onset heart failure (2004-2010).

The researchers found that from examination 1 to examination 2, the prevalence of diastolic dysfunction of any degree increased from 23.8 percent to 39.2 percent. Moderate or severe diastolic dysfunction increased from 6.4 percent to 16.0 percent. Over 4 years, 23.4 percent of participants experienced worsening of diastolic function, 67.8 percent remained unchanged, and 8.8 percent experienced improved diastolic function. Age was predictive of the development of diastolic dysfunction dysfunction, especially ages 65 years or older.

In the analysis of the diastolic function of healthy participants (without hypertension, diabetes, coronary artery disease, heart failure, or use of cardiovascular medications), incidence of diastolic dysfunction of any degree increased from 11.3 percent at examination 1 to 29.8 percent at examination 2. Among 423 healthy participants, 19.9 percent showed worsening diastolic function, 75.2 percent remained the same, and 5.0 percent improved.

During 6.3 years of additional follow-up, 81 participants developed heart failure. Age 65 years or older was the strongest predictor of heart failure. The authors found that persistent or worsening diastolic dysfunction was associated with heart failure. "Cumulative heart failure incidence was 2.6 percent in participants whose diastolic function remained normal or normalized between examinations; 7.8 percent in those with persistent, or progression to mild diastolic dysfunction; and 12.2 percent in those with persistent, or progression to moderate or severe diastolic dysfunction."



"Longitudinal evaluation of <u>participants</u> in the population-based OCHFS cohort reveals that left ventricular diastolic dysfunction is highly prevalent, tends to worsen over time, and is associated with advancing age. Worsening diastolic function can be detected even in apparently healthy persons. Although confirmation in other studies would be helpful, our data suggest that persistence or progression of <u>diastolic</u> <u>dysfunction</u> is a risk factor for <u>heart failure</u> in elderly persons," the researchers conclude.

More information: JAMA. 2011;306[8]:856-863.

Provided by JAMA and Archives Journals

Citation: Diastolic dysfunction appears to worsen over time; associated with increased risk of heart failure (2011, August 23) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2011-08-diastolic-dysfunction-worsen-heart-failure.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.