

Study finds flexible-fiber CO2 laser safe in endoscopic endonasal transsphenoidal surgery

August 16 2011

A new study has shown the Flexible-Fiber CO2 laser to safely cut and coagulate during endoscopic assisted transsphenoidal craniotomies (TSC) without the line of sight problems encountered with conventional CO2 lasers. The findings, reported by researchers from Boston Medical Center (BMC) and Boston University School of Medicine (BUSM), appear in the July/August issue of *World Neurosurgery*. This is the first study to report the utility of flexible-fiber CO2 laser-assisted ablation in TSC.

The implementation of [endoscopy](#) in transsphenoidal surgery for pituitary tumors and other sellar lesions has been one of the most recent significant advances for the treatment of these tumors. This is bolstered by the demonstration of its safety and relative advantages over traditional microscope-based approaches. The long wavelength of the carbon dioxide (CO2) laser has been demonstrated to have high absorption in tissue and water, with rapid conversion of [light energy](#) into heat within a small volume of tissue. This property makes the CO2 laser an excellent cutting and [ablation](#) tool because it causes minimal thermal damage to adjacent tissue.

The researchers performed a retrospective chart review on 16 patients who underwent flexible-fiber CO2 laser-assisted endoscopic ESS. All tumor pathology types were considered. Results were assessed based on hormone status, [tumor size](#), pathology, complications and resection rates.

Pituitary lesions with an average size of 22.7 mm were identified by radiographic and pathologic criteria. The researchers found surgery had a low rate of complication, (three patients) and no laser-related complications were encountered.

According to the authors, because of the small working space, the propensity for bleeding, and the amount of soft tissue and osteogenic dissection involved, the use of flexible-fiber CO2 lasers in ESS is ideal.

Because this is a feasibility and safety study, the authors cannot draw any conclusions regarding outcomes compared with traditional endoscopic transsphenoidal methods, including any difference in time required for laser surgery. However, as their experience has increased they have seen improved time efficiency with regard to laser set up and use.

"The flexible-fiber CO2 laser tool described and utilized in this report is ideally suited for endoscopic assisted TSC because of the narrow corridors and difficulty in using conventional cutting and coagulating instruments," said study co-author author Anand K. Devaiah, MD, FACS, an otolaryngologist at BMC and associate professor of otolaryngology and neurological surgery at BUSM. Devaiah has worked extensively with this system in pioneering applications for minimally invasive surgery, and works with the CO2 laser fiber company, OmniGuide, as a consultant. "Because most pituitary tumors are soft, the CO2 [laser](#) was not used extensively for tumor resection in this series, but we have found it very helpful in removing recurrent pituitary adenomas where tumor consistency is often more fibrous and dural scarring may be extensive," added Devaiah.

"Further study of this promising surgical tool is being investigated in other areas of neurosurgery including tumors of the brains and spinal cord," said senior author Lawrence Chin, MD, chief of neurosurgery at BMC and professor and chairman of neurosurgery at BUSM.

Provided by Boston University Medical Center

Citation: Study finds flexible-fiber CO2 laser safe in endoscopic endonasal transsphenoidal surgery (2011, August 16) retrieved 25 April 2024 from <https://medicalxpress.com/news/2011-08-flexible-fiber-co2-laser-safe-endoscopic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.