

Some hospitals better than others in selecting patients to undergo cardiac catheterization

August 9 2011

(Medical Xpress) -- Hospitals vary markedly when it comes to the rate at which diagnostic coronary angiography or catheterization – an invasive procedure that allows doctors to see the vessels and arteries leading to the heart – actually finds obstructive coronary artery disease (CAD) in people without known heart disease.

In fact, while some U.S. hospitals report that 100 percent of patients undergoing this procedure were found to have [CAD](#), others had rates as low as 23 percent, meaning the majority of patients selected for elective catheterization did not have blockages, according to a new study published in the August 16, 2011, issue of the *Journal of the American College of Cardiology*. Researchers say these findings warrant further efforts to improve the patient selection and decision-making processes used by institutions to limit exposure to costly, invasive procedures when not needed.

"This procedure has associated costs and it's not without risk," said Pamela S. Douglas, MD, Ursula Geller Professor of Cardiovascular Research, Duke Clinical Research Institute, Durham, NC, and lead author of the study. "This study is an important step in assessing quality of care and is integral to efforts to improve it. Our findings indicate that there may be an opportunity to increase the likelihood of finding CAD at catheterization, and perhaps reduce the number of procedures that don't find disease."

The study – a follow up to the team's 2010 study – is the first to

investigate the degree to which hospitals differ in the rate of discovering CAD with coronary angiography and the factors that might predict this. Hospital-level variability appears to be predictable based on differing patterns of patient selection and pre-catheterization evaluation, testing and treatment.

"Because hospitals maintained a similar rate of finding disease relative to other hospitals year in and year out, and these rates are also related to patient characteristics, it suggests that decision-making processes and clinical practice patterns are a highly influential factor guiding the use of diagnostic [coronary angiography](#) and could be a target for quality improvement efforts," explained Dr. Douglas.

While the study was not able to capture all of the reasons for performing angiography, its findings suggest that some procedures might be avoided if decisions about whether to perform cardiac catheterization in the hospitals with the lowest rates of finding CAD were more similar to those hospitals with the highest rates.

Hospitals with lower rates of finding CAD by catheterization were more likely to perform the procedures on younger patients, women, those with a lower likelihood of disease (as indicated by a low Framingham or Diamond and Forrester risk score), who presented with atypical symptoms and who had negative non-invasive diagnostic testing done prior to angiography. Not surprisingly, patients at higher risk of being diagnosed with obstructive coronary artery disease were more likely to have hypertension, diabetes and other expected risk factors. In general, the size and type of hospital (e.g., teaching [hospital](#), public vs. private) and setting did not change the findings.

Researchers identified 565,504 patients without known heart disease who underwent elective cardiac catheterization at 691 hospitals nationwide over a three-year-period to evaluate the rate of finding

obstructive CAD. Authors defined CAD as any major epicardial vessel stenosis at 50 percent or higher, but similar patterns remained even when alternate definitions were applied.

"The decision to perform catheterization should be selective and ideally limited to patients with moderate to high pre-test probability for CAD," said Dr. Douglas, who also cautions that there are other, valid reasons to do a catheterization besides the expectation of finding obstructive CAD, such as patient reassurance.

To provide additional guidance to cardiologists, ACC is currently working on appropriate use criteria for ordering invasive diagnostic coronary angiographies with release expected this winter. In the meantime, Dr. Douglas advocates for closer consideration of those clinical factors known to be most strongly associated with CAD, including advancing age, risk factors and typical symptoms. A careful assessment of patients' risk and presenting symptoms, as well as results of any stress and other non-invasive tests should be considered.

Provided by American College of Cardiology

Citation: Some hospitals better than others in selecting patients to undergo cardiac catheterization (2011, August 9) retrieved 24 April 2024 from <https://medicalxpress.com/news/2011-08-hospitals-patients-cardiac-catheterization.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.